

## Technical Data Sheet


## Biotin Rat Anti-Mouse CD195

## Product Information

Material Number:	559922
Alternate Name:	CCR5
Size:	0.5 mg
Concentration:	0.5 mg/ml
Clone:	C34-3448
Immunogen:	Mouse CCR5 aa. 9-30
Isotype:	Rat IgG2c, $\kappa$
Reactivity:	QC Testing: Mouse
Storage Buffer:	Aqueous buffered solution containing $\leq 0.09\%$ sodium azide.

## Description

The C34-3448 antibody reacts with mouse chemokine receptor CD195 (also known as CCR5), a member of the  $\beta$ -chemokine receptor family. CD195 regulates lymphocyte chemotaxis activation and transendothelial migration during inflammation. It signals a response to at least three chemokines: RANTES and macrophage inflammatory protein-1 (MIP-1) $\alpha$  and  $\beta$ . In the mouse system, the gene for CD195 has been mapped to chromosome 9. CD195 mRNA is expressed in heart, spleen and liver tissues, and by macrophages and T-lymphocytes. The immunogen used to generate the C34-3448 hybridoma was a KLH-conjugated peptide consisting of amino acids 9-30 of mouse CD195 (CCR5).



**Expression of mouse CD195 by stimulated mouse splenocytes.** BALB/c splenocytes were stimulated for 24 hours with ConA (10  $\mu$ g/ml, final concentration; Sigma). The splenocytes were cultured for additional 5 days in the presence of mouse IL-2 (100 U/ml, final concentration; Cat. No. 550069). The splenocytes were stained with biotinylated rat anti-mouse CD195 antibody (C34-3448) and followed by Streptavidin-Phycoerythrin (Cat. No. 554061) following the BD Pharmingen staining protocol (see Figure, Left panel). The data reflects gating on lymphocytes, based on forward and side scattered light signals. The level of nonspecific staining was assessed using the biotinylated rat IgG2c isotype control (A23-1; Cat. No. 553983; Right panel). The quadrant markers for the bivariate dot plots were set based on the autofluorescence control and isotype control.

## Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with biotin under optimum conditions, and unreacted biotin was removed.

Store undiluted at 4° C and protected from prolonged exposure to light. Do not freeze.

## BD Biosciences

[www.bdbiosciences.com](http://www.bdbiosciences.com)

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country-specific contact information, visit [www.bdbiosciences.com/how\\_to\\_order/](http://www.bdbiosciences.com/how_to_order/)

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2007 BD



## Application Notes

### Application

Flow cytometry	Routinely Tested
----------------	------------------

### Recommended Assay Procedure:

**Immunofluorescent Staining and Flow Cytometry Analysis:** The biotin-conjugated C34- 3448 antibody can be used for multicolor immunofluorescent staining and flow cytometric analyses to identify and enumerate CD195-expressing cells within mixed cell populations. For optimal immunofluorescent staining with flow cytometric analysis, the antibody should be titrated ( $\leq 0.25$   $\mu$ g mAb/million cells). An appropriate isotype control for this antibody is Cat. No. 553983. For specific methodology, please visit the protocols section online or the chapter on intracellular staining and flow cytometry in the Immune Function Handbook. Both of these resources are on our web site, [www.bdbiosciences.com](http://www.bdbiosciences.com).

### Suggested Companion Products

Catalog Number	Name	Size	Clone
550069	Recombinant mouse IL-2	20 $\mu$ g	(none)
554061	PE Streptavidin	0.5 mg	(none)
553983	Biotin Rat IgG2c, $\kappa$ Isotype Control	0.25 mg	A23-1

### Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to [www.bdbiosciences.com/pharming/en/protocols](http://www.bdbiosciences.com/pharming/en/protocols) for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

### References

Boring L, Gosling J, Montecarlo FS, Lulis AJ, Tsou CL, Charo IF. Molecular cloning and functional expression of murine JE (monocyte chemoattractant protein 1) and murine macrophage inflammatory protein 1alpha receptors: evidence for two closely linked C-C chemokine receptors on chromosome 9. *J Biol Chem.* 1996; 271(13):7551-7558.(Biology)

Meyer A, Coyle AJ, Proudfoot AE, Wells TN, Power CA. Cloning and characterization of a novel murine macrophage inflammatory protein-1 alpha receptor. *J Biol Chem.* 1996; 271(24):14445-14451.(Biology)

Napolitano M, Seamon KB, Leonard WJ. Identification of cell surface receptors for the Act-2 cytokine. *J Exp Med.* 1990; 172(1):285-289.(Biology)