BD™ Cytometric Bead Array (CBA)



Technical Data Sheet Human Soluble CD121a (IL-1 RI) Flex Set

Product Information

- Material Number: Size: Bead Position: Assay Range: Reactivity:
- 560276 100 Tests B6 40-10,000 pg/mL QC Testing: Human

Component Description: Component Mat. No: Component Storage Buffer:

Component Description: Component Mat. No: Component Storage Buffer:

Component Description: Component Mat. No: Component Storage Buffer: 51-9005889
Lyophilized in an aqueous buffered solution containing BSA and ProClin[™] 150.
Human Soluble CD121a (IL-1 RI) PE Detection Reagent 51-9005891

Human Soluble CD121a (IL-1 RI) Standard

Aqueous buffered solution containing BSA and $\leq 0.09\%$ sodium azide.

Human Soluble CD121a (IL-1 RI) Capture Bead B6 51-9005893 Aqueous buffered solution containing fetal bovine serum and ≤0.09% sodium azide.




Figure 1. Example BD CBA Human Soluble CD121a (IL-1 RI) Flex Set standard curve. Data acquired on a BD FACSArray bioanalyzer and analyzed using FCAP Array Software.

Description

The BDTM Human Soluble CD121a (IL-1 RI) Flex Set is a bead-based immunoassay capable of measuring soluble human soluble sCD121a, also known as IL-1 Receptor type-1, in serum, plasma, and cell culture supernatant samples. Human reactivity was determined by testing samples with the BD CBA Human Soluble CD121a (IL-1 RI) Flex Set. The biology and function of sCD121a (IL-1 Receptor type-1) has been extensively reviewed in the literature.

Preparation and Storage

This BD[™] CBA Flex Set contains one vial each of Capture Bead and PE Detection Reagent and two vials of Standard. The Capture Bead and PE Detection Reagent components of this flex set have been formulated to a 50x concentration to ensure product performance when multiplexed. The Standard component is lyophilized and should be transferred to a 15 mL polypropylene tube for reconstituted standard, do not reconstituted in 4.0 mL Assay Diluent, the standard has a protein concentration of 10,000 pg/mL. Discard unused reconstituted standard, do not store or reuse. Store lyophilized standard and other components at 4°C. Protect Capture Beads and the PE Detection Reagent from prolonged exposure to light.

Application Notes

Recommended Assay Procedure: The BD CBA Human Soluble CD121a (IL-1 RI) Flex Set must be used in conjunction with a BD CBA Human Soluble Protein Master Buffer Kit (Cat. No. 558264, 100 tests, or 558265, 500 tests), a flow cytometer, and FCAP Array[™] Software. Detailed instructions on the use of this product can be found in the manual for the BD CBA Human Soluble Protein Master Buffer Kit. When following the directions in the Master Buffer Kit, the top standard point for the BD CBA Human Soluble CD121a (IL-1 RI) Flex Set will be 10,000 pg/mL. An example standard curve is shown in Figure 1.

The BD CBA Human Soluble CD121a (IL-1 RI) Flex Set should not be used in the same assay well with any non-BD CBA Human Soluble

BD Biosciences

bdbiosciences.	om				
United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbear
877.232.8995	800.268.5430	32.2.400.98.95	0120.8555.90	65.6861.0633	55.11.5185.9995
For country cor	ntact informatio	on, visit bdbiosci e	ences.com/conta	ct	
of any patents. BL	D Biosciences will n	ot be held responsik	ble for patent infrin	gement or other vio	e the above product in violation
use of our product	ts. Purchase does n	tot include or carry a	any right to resell or	r transfer this produ	lations that may occur with the
product or as a co	mponent of anoth	er product. Any use	of this product oth	er than the permitte	ct either as a stand-alone
written authoriza	tion of Becton, Dic	kinson and Compan	y is stictly prohibite	d.	ed use without the express
For Research Use	Only. Not for use ir	diagnostic or thera	apeutic procedures.	Not for resale.	in and Company. © 2014 BD



Receptor or CD Marker Flex Set reagents (such as BD CBA Mouse Soluble Protein or Cell Signaling Flex Sets) **nor with any BD CBA Human Soluble Protein Flex Set reagents**. For an updated assay compatibility chart for the BD CBA Human Soluble Protein Flex Sets, please refer to the BD CBA Flex Set System homepage at http://www.bdbiosciences.com/cbasetup.

Performance

Limit of Detection: The theoretical limit of detection is 2.3 pg/mL and was determined by evaluating the estimated result of the average MFI of the negative control (0 pg/mL, n=30) + 2 standard deviations.

	Inter-Assay Reproducibility			Intra-Assay Reproducibility			
Specificity		Mean (pg/ml)	Standard Deviation	%CV	Mean (pg/ml)	Standard Deviation	%CV
	Sample 1	150.9	11.4	8%	158.8	5.1	3%
Human sCD121a Sample 2		592.3	22.6	4%	599.4	25.9	4%
(IL-1 RI)	Sample 3	2573.4	204.8	8%	2471.3	138.0	6%

Reproducibility: The intra-assay and inter-assay reproducibility were determined for the BD CBA Human Soluble CD121a (IL-1 RI) Flex Set by evaluating ten replicates of three different sample levels (intra-assay) and three replicates of three different sample levels from four separate experiments (inter-assay).

	Cell Culture Supernatant		Ser	um	Plasma	
Specificity	Average % Recovery	Range	Average % Recovery	Range	Average % Recovery	Range
Human sCD121a (IL-1 RI)	99%	92 - 105%	87%	85 - 88%	73%	72 - 75%

Recovery: Cell culture supernatant, serum, or EDTA-treated plasma were spiked with three different levels of protein. The spiked samples were assayed and the results were compared with expected values. Serum and plasma samples were diluted 1:4 before the protein was spiked into each. Serum is a pool of 800 - 1000 donors and the plasma was pooled from at least 20 donors.

Comula Dilution	Cell Culture Supernatant		Ser	um	Plasma		
Sample Dilution	Detected (pg/mL)	% of Expected	Detected (pg/mL)	% of Expected	Detected (pg/mL)	% of Expected	
Spiked sample	5195.5	100%	4624.5	100%	4324.0	100%	
1:2	2404.1	93%	2282.3	99%	2210.8	102%	
1:4	1256.5	97%	1223.3	106%	1207.5	112%	

Linearity: Cell culture supernatant, 1:4 diluted serum, or 1:4 diluted EDTA-treated plasma were spiked with protein and serially diluted. The diluted samples were assayed and the results were compared with the original spiked sample.

Product Notices

1. ProClin is a trademark of Rohm and Haas Company.

- 2. Source of all serum proteins is from USDA inspected abattoirs located in the United States.
- 3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
- Warning: CBA lyophilized standard contains 0.02% (w/w) of a CMIT/MIT mixture (3:1), which is a mixture of: 5-chloro-2-methyl-4-isothiazolin-3-one [EC No 247-500-7] and 2-methyl-4-isothiazolin-3-one [EC No 220-239-6] (3:1). Hazard statement: May cause an allergic skin reaction.

Precautionary statements: Wear protective gloves/eye protection. Wear protective clothing. Avoid breathing mist/vapours/spray. If skin irritation or rash occurs: Get medical advice/attention. IF ON SKIN: Wash with plenty of water. Dispose of contents/container in accordance with local/regional/national/international regulations.

BD Biosciences bdbiosciences.com								
babiosciences.	lom							
United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean			
877.232.8995	800.268.5430	32.2.400.98.95	0120.8555.90	65.6861.0633	55.11.5185.9995			
For country co	ntact informatio	on, visit bdbiosci e	ences.com/conta	ct				
of any patents. BL use of our produc	D Biosciences will n ts. Purchase does n	ot be held responsil ot include or carry a	ole for patent infrin any right to resell or	gement or other vio r transfer this produ	e the above product in violation lations that may occur with the ct either as a stand-alone ed use without the express			



written authorization of Becton, Dickinson and Company is stictly prohibited. For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2014 BD