

Technical Data Sheet

PE Mouse anti-Pyk2 (pY402)


Product Information

| | |
|-------------------------|--|
| Material Number: | 560255 |
| Alternate Name: | Protein tyrosine kinase 2 β , FAK2, FADK 2, CAK β , CADTK, RAFTK |
| Size: | 50 tests |
| Vol. per Test: | 20 μ l |
| Clone: | L68-1256.272 |
| Immunogen: | Phosphorylated Human Pyk2 Peptide |
| Isotype: | Mouse (BALB/c) IgG2b, κ |
| Reactivity: | QC Testing: Human Predicted: Mouse, Rat |
| Storage Buffer: | Aqueous buffered solution containing BSA and \leq 0.09% sodium azide. |

Description

Proline-rich tyrosine kinase 2 (Pyk2) is a member of the FAK subfamily of tyrosine protein kinases. Pyk2 responds to a wide variety of inflammatory and stress stimuli, such as cytokines, ligation of integrins or antigen receptors, mechanical strain, and cellular depolarization. A common characteristic of most Pyk2 stimuli is an increase in cytosolic free calcium. Pyk2 is involved in signaling pathways that regulate vital cellular functions in hematopoietic, epithelial, neuronal, endothelial, and other cell types. Activation of Pyk2 initiates its autophosphorylation at tyrosine 402 (Y402), followed by Src-mediated phosphorylation of other Pyk2 tyrosine sites, which enhances its activity and leads to downstream signal transduction through the MAPK and JNK pathways.

The L68-1256.272 monoclonal antibody recognizes the phosphorylated Y402 of activated Pyk2.



Analysis of Pyk2 (pY402) in human peripheral blood lymphocytes. Human peripheral blood mononuclear cells (PBMC) were either untreated (left panel) or stimulated by cross-linking of CD3 and CD28 with NA/LE Mouse anti-Human CD3 mAb UCHT1 (Cat. No. 555329) and NA/LE Mouse anti-Human CD28 mAb CD28.2 (Cat. No. 555725) on ice for 15 minutes followed by Purified Goat anti-Mouse Ig (Cat. No. 553998) on ice for 15 minutes, and then allowed to undergo phosphorylation at 37°C for 5 minutes (right panel). The cells were fixed (BD Cytotfix™ buffer, Cat. No. 554655) for 10 minutes at 37°C, permeabilized (BD™ Phosflow Perm Buffer III, Cat. No. 558050) on ice for 30 minutes, blocked with normal mouse immunoglobulin, and then stained with PE Mouse anti-Pyk2 (pY402) and APC Mouse anti-human CD3 mAb SK7 (Cat. No. 340440). For data analysis, lymphocytes were selected by their scatter profile. The data demonstrates that the upregulated phosphorylation of Pyk2 was restricted to the CD3-positive T lymphocytes under these stimulation conditions. Flow cytometry was performed on a BD™ FACSCalibur flow cytometry system.

The specificity of mAb L68-1256.272 was confirmed by western blot analysis using unconjugated antibody on lysates from PBMC that were untreated (lane 1) or stimulated by cross-linking of CD3 and CD28 (lane 2). Pyk2 (pY402) is identified as a band of 116 kDa, with increased intensity in the stimulated cells.

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with R-PE under optimum conditions, and unconjugated antibody and free PE were removed.

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

BD Biosciences

bdbiosciences.com

| | | | | | |
|---------------|--------------|---------------|--------------|--------------|-------------------------|
| United States | Canada | Europe | Japan | Asia Pacific | Latin America/Caribbean |
| 877.232.8995 | 888.268.5430 | 32.53.720.550 | 0120.8555.90 | 65.6861.0633 | 0800.771.7157 |

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2011 BD



The purified or conjugated mAb was characterized by flow cytometry (Flow) and western blot (WB) using these model systems:

| Method | Species | Cells | Treatment | Fixation | Perm buffer | Result |
|--------|---------|--------|-----------------------|------------------|--------------------|----------------------|
| Flow | Human | Jurkat | Hydrogen peroxide | Cytofix or Fix I | Perm III | Induction observed |
| | Human | PBMC | CD3/CD28 crosslinking | Cytofix or Fix I | Perm I, II, or III | Induction observed |
| WB | Human | Jurkat | Hydrogen peroxide | | | 116-kDa band induced |
| | Human | Jurkat | anti-CD3 | | | Unsatisfactory |
| | Human | PBMC | CD3/CD28 crosslinking | | | 116-kDa band induced |

Application Notes

Application

| | |
|---|------------------|
| Intracellular staining (flow cytometry) | Routinely Tested |
|---|------------------|

Recommended Assay Procedure:

Either BD Cytofix™ fixation buffer or BD™ Phosflow Fix Buffer I may be used for cell fixation. Any of the three BD™ Phosflow permeabilization buffers may be used.

Suggested Companion Products

| Catalog Number | Name | Size | Clone |
|----------------|--------------------|--------|--------|
| 554655 | Fixation Buffer | 100 ml | (none) |
| 557870 | Fix Buffer I | 250 ml | (none) |
| 557885 | Perm/Wash Buffer I | 125 ml | (none) |
| 558052 | Perm Buffer II | 125 ml | (none) |
| 558050 | Perm Buffer III | 125 ml | (none) |

Product Notices

1. This reagent has been pre-diluted for use at the recommended Volume per Test. We typically use 1×10^6 cells in a 100- μ l experimental sample (a test).
2. Please refer to www.bdbiosciences.com/pharming/en/protocols for technical protocols.
3. For fluorochrome spectra and suitable instrument settings, please refer to our Fluorochrome Web Page at www.bdbiosciences.com/colors.
4. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
5. Source of all serum proteins is from USDA inspected abattoirs located in the United States.

References

- Anand AR, Cucchiari M, Terwilliger EF, Ganju RK. The tyrosine kinase Pyk2 mediates lipopolysaccharide-induced IL-8 expression in human endothelial cells. *J Immunol.* 2008; 180:5636-5644. (Biology)
- Avraham H, Park S-Y, Schinkmann K, Avraham S. RAFTK/Pyk2-mediated cellular signalling. *Cell Signal.* 2000; 12(3):123-133. (Biology)
- Faure C, Corvol J-C, Toutant M, et al. Calcineurin is essential for depolarization-induced nuclear translocation and tyrosine phosphorylation of PYK2 in neurons. *J Cell Sci.* 2007; 120:3034-3044. (Biology)
- Gluck SL. Acid sensing in renal epithelial cells. *J Clin Invest.* 2004; 114(12):1696-1699. (Biology)
- Orr AW, Murphy-Ullrich EM. Regulation of endothelial cell function by FAK and Pyk2. *Front Biosci.* 2004; 9:1254-1266. (Biology)
- Park, S-Y, Avraham HK, Avraham S. RAFTK/Pyk2 activation is mediated by trans-acting autophosphorylation in a Src-independent manner. *J Biol Chem.* 2004; 279(32):33315-33322. (Biology)