


Technical Data Sheet

Alexa Fluor® 647 Mouse anti-β-Tubulin, Class III**Product Information**

| | |
|-------------------------|--|
| Material Number: | 560394 |
| Alternate Name: | tubulin, beta 3; MC1R; TUBB3; TUBB4; tubulin, beta-4 |
| Size: | 50 tests |
| Vol. per Test: | 20 µl |
| Clone: | TUJ1 |
| Immunogen: | Rat brain microtubules |
| Isotype: | Mouse IgG2a |
| Reactivity: | QC Testing: Human Reported Reactivity: Rat |
| Storage Buffer: | Aqueous buffered solution containing BSA, protein stabilizer, and ≤0.09% sodium azide. |

Description

Microtubules are formed by the self assembly of tubulin and are one of the major components of the eukaryotic cytoskeleton. The two main tubulin isoforms, α - and β -tubulin, are usually products of separate genes. The β -tubulin family includes six expressed genes that produce the polypeptide isoforms known as Classes I through VI, each of which have a distinct pattern of expression. Class III β -tubulin is found in neurons and mammalian testis cells and is widely used as a neuronal marker in developmental neurobiology, neoplasia, and stem cell research. Class III β -tubulin expression in neuronal and neuroblastic tumors is differentiation dependent, and its expression in certain non-neuronal neoplasms has been associated with poor prognosis and/or resistance to chemotherapy.



Analysis of Alexa Fluor® 647 Anti-Human Class III tubulin on Neurons. H9 human embryonic stem (ES) cells (WiCell, Madison, WI) were differentiated into Neural Precursor cells (NPCs) and grown for 4 passages before differentiating into neurons and glia for 12 days. The cells were fixed with BD Cytotfix™ Fixation Buffer (Cat. No. 554655) for 20 minutes at room temperature, permeabilized with BD Phosflow™ Perm/Wash Buffer I (Cat. No. 557885), and then stained with either Alexa Fluor® 647 Mouse anti-β-Tubulin, Class III (Cat. No. 560394; Right Histogram) or Alexa Fluor® 647 Mouse IgG2a Isotype Control (Cat. No. 558053; Left Histogram). This antibody also works with BD Phosflow™ Perm Buffer II and III. Flow cytometry was performed on a BD FACSCanto™ II flow cytometry system.

Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated to Alexa Fluor® 647 under optimum conditions, and unreacted Alexa Fluor® 647 was removed.

Application Notes**Application**

| | |
|---|------------------|
| Intracellular staining (flow cytometry) | Routinely Tested |
| Bioimaging | Routinely Tested |

Suggested Companion Products

| Catalog Number | Name | Size | Clone |
|----------------|---|----------|----------|
| 554655 | Fixation Buffer | 100 ml | (none) |
| 557885 | Perm/Wash Buffer I | 125 ml | (none) |
| 558053 | Alexa Fluor® 647 Mouse IgG2a, κ Isotype Control | 50 tests | MOPC-173 |
| 554656 | Stain Buffer (FBS) | 500 ml | (none) |

BD Biosciences

bdbiosciences.com

United States 877.232.8995 Canada 800.268.5430 Europe 32.2.400.98.95 Japan 0120.8555.90 Asia Pacific 65.6861.0633 Latin America/Caribbean 55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2014 BD



Product Notices

1. Please refer to www.bdbiosciences.com/pharming/protocols for technical protocols.
2. This reagent has been pre-diluted for use at the recommended Volume per Test. We typically use 1×10^6 cells in a 100- μ l experimental sample (a test).
3. Alexa Fluor® 647 fluorochrome emission is collected at the same instrument settings as for allophycocyanin (APC).
4. Alexa Fluor® is a registered trademark of Molecular Probes, Inc., Eugene, OR.
5. The Alexa Fluor®, Pacific Blue™, and Cascade Blue® dye antibody conjugates in this product are sold under license from Molecular Probes, Inc. for research use only, excluding use in combination with microarrays, or as analyte specific reagents. The Alexa Fluor® dyes (except for Alexa Fluor® 430), Pacific Blue™ dye, and Cascade Blue® dye are covered by pending and issued patents.
6. For fluorochrome spectra and suitable instrument settings, please refer to our Multicolor Flow Cytometry web page at www.bdbiosciences.com/colors.
7. Source of all serum proteins is from USDA inspected abattoirs located in the United States.
8. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
9. An isotype control should be used at the same concentration as the antibody of interest.
10. This reagent has been pre-diluted for use at the recommended Volume per Test when following the Recommended Assay Procedure. A Test is typically ~10,000 cells cultured in a well of a 96-well imaging plate.

References

Geisert EE Jr, Frankfurter A. The neuronal response to injury as visualized by immunostaining of class III beta-tubulin in the rat. *Neurosci Lett.* 1989; 102(2-3):137-141. (Clone-specific: Immunohistochemistry)

Katsetos CD, Del Valle L, Geddes JF et al. Aberrant localization of the neuronal class III beta-tubulin in astrocytomas. *Arch Pathol Lab Med.* 2001; 125(5):613-624. (Clone-specific: Immunocytochemistry (cytospins))

Mozzetti S, Ferlini C, Concolino P et al. Class III beta-tubulin overexpression is a prominent mechanism of paclitaxel resistance in ovarian cancer patients. *Clin Cancer Res.* 2005; 11(1):298-305. (Clone-specific: Immunohistochemistry)

BD Biosciences

bdbiosciences.com

| United States | Canada | Europe | Japan | Asia Pacific | Latin America/Caribbean |
|---------------|--------------|----------------|--------------|--------------|-------------------------|
| 877.232.8995 | 800.268.5430 | 32.2.400.98.95 | 0120.8555.90 | 65.6861.0633 | 55.11.5185.9995 |

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2014 BD

