

Technical Data Sheet

Alexa Fluor® 488 Mouse anti-Elk-1 (pS383)


Product Information

Material Number:	560417
Alternate Name:	ELK1
Size:	50 tests
Vol. per Test:	20 µl
Clone:	M21-1721
Immunogen:	Phosphorylated Human Elk-1 Peptide
Isotype:	Mouse (BALB/c) IgG1, κ
Reactivity:	QC Testing: Human Predicted due to immunogen sequence identity: Mouse, Rat
Storage Buffer:	Aqueous buffered solution containing BSA, protein stabilizer, and ≤0.09% sodium azide.


Description

Elk-1 is a 62-kDa member of the Ets oncogene family of transcription factors and a member of the subfamily of ternary complex factors (TCF). Ets proteins mediate a variety of gene activities in response to serum and growth factors. Proteins in the TCF subfamily form a ternary complex by binding to the Serum Response Element (SRE) in conjunction with a dimer of Serum Response Factors (SRF). Elk-1 is phosphorylated by mitogen-activated protein (MAP) kinase pathways in vivo at a cluster of S/T motifs at its carboxy-terminus. Phosphorylation at these sites, particularly at serine 383 (S383), is critical for Elk-1 transcriptional activation. Studies have shown that Elk-1 is a direct target of activated MAP kinase, and that it is also a target of the stress-activated kinase SAPK/JNK.

The M21-1721 monoclonal antibody recognizes the phosphorylated S383 of activated Elk-1. The orthologous phosphorylation sites of mouse and rat Elk-1 are S384 and S382, respectively.



Analysis Elk-1 (pS383) in activated human peripheral blood mononuclear cells (PBMC). PBMC were isolated by density gradient centrifugation (Ficoll-Paque™ PLUS, Cat. No. 17-1440-02), and cultured with 20 µg/ml PHA-P (Sigma-Aldrich, Cat. No. L1668) for 3 days, then either left untreated (open histogram) or treated with PMA at 50 nM/10⁶ cells for 15 minutes (shaded histogram; Sigma-Aldrich, Cat. No. P8139). Cells were then fixed in BD Cytotfix™ buffer (Cat. No. 554655) at 37°C for 10 minutes, then permeabilized with BD Phosflow™ Perm Buffer III (Cat. No. 558050) on ice for at least 30 minutes, and then stained with Alexa Fluor® 488 Mouse anti-Elk-1 (pS383). Flow cytometry was performed on a BD FACSCalibur™ flow cytometry system.



Western blot analysis of Elk-1 (pS383). The specificity of mAb M21-1721 was confirmed by western blot analysis using unconjugated Mouse anti-Elk-1 (pS383) antibody on lysates from PBMC that were cultured with PHA for 3 days (lane 1) or PBMC that were cultured with PHA for 3 days and then treated with PMA for 15 minutes (lane 2). Elk-1 (pS383) is identified as a band of 62 kDa in the PMA-treated cells.

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	800.979.9408	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD

Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated to Alexa Fluor® 488 under optimum conditions, and unreacted Alexa Fluor® 488 was removed.

The purified or conjugated mAb was characterized by flow cytometry (Flow) and western blot (WB) using these model systems:

Method	Species	Cells	Treatment	Fixation	Perm buffer	Result
Flow	Human	PHA-stimulated PBMC	PMA	Cytofix	Perm I, II, or III	Upregulated expression
WB	Human	PHA-stimulated PBMC	PMA			62-kDa band induced
		HeLa S3	PMA			62-kDa band induced

Application Notes

Application

Intracellular staining (flow cytometry)	Routinely Tested
---	------------------

Recommended Assay Procedure:

This antibody conjugate is suitable for intracellular staining of human peripheral blood mononuclear cells using BD Cytotfix™ Fixation Buffer.

Any of the three BD Phosflow™ permeabilization buffers may be used.

Suggested Companion Products

Catalog Number	Name	Size	Clone
554655	Fixation Buffer	100 ml	(none)
557885	Perm/Wash Buffer I	125 ml	(none)
558052	Perm Buffer II	125 ml	(none)
558050	Perm Buffer III	125 ml	(none)

Product Notices

1. This reagent has been pre-diluted for use at the recommended Volume per Test. We typically use 1×10^6 cells in a 100- μ l experimental sample (a test).
2. Please refer to www.bdbiosciences.com/pharming/protocols for technical protocols.
3. Alexa Fluor® 488 fluorochrome emission is collected at the same instrument settings as for fluorescein isothiocyanate (FITC).
4. For fluorochrome spectra and suitable instrument settings, please refer to our Multicolor Flow Cytometry web page at www.bdbiosciences.com/colors.
5. Alexa Fluor® is a registered trademark of Molecular Probes, Inc., Eugene, OR.
6. The Alexa Fluor®, Pacific Blue™, and Cascade Blue® dye antibody conjugates in this product are sold under license from Molecular Probes, Inc. for research use only, excluding use in combination with microarrays, or as analyte specific reagents. The Alexa Fluor® dyes (except for Alexa Fluor® 430), Pacific Blue™ dye, and Cascade Blue® dye are covered by pending and issued patents.
7. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
8. Source of all serum proteins is from USDA inspected abattoirs located in the United States.
9. Ficoll-Paque is a trademark of Amersham Biosciences Limited.
10. All other brands are trademarks of their respective owners.

References

Bardwell AJ, Abdollahi M, Bardwell L. Docking sites on mitogen-activated protein kinase (MAPK) kinases, MAPK phosphatases and the Elk-1 transcription factor compete for MAPK binding and are crucial for enzymic activity. *Biochem J.* 2003; 370:1077-1085. (Biology)

Cavigelli M, Dolfi F, Claret FX, Karin M. Induction of c-fos expression through JNK-mediated TCF/Elk-1 phosphorylation. *EMBO J.* 1995; 14(23):5957-5964. (Biology)

Espanel X, Wälchli S, Rüdke T et al. Mapping of synergistic components of weakly interacting protein-protein motifs using arrays of paired peptides. *J Biol Chem.* 2003; 278(17):15162-15167. (Biology: Bioassay)

Sugimoto T, Stewart S, Guan KL. The calcium/calmodulin-dependent protein phosphatase calcineurin is the major Elk-1 phosphatase. *J Biol Chem.* 1997; 272(47):29415-29418. (Biology)

Tian J, Karin M. Stimulation of Elk1 transcriptional activity by mitogen-activated protein kinases is negatively regulated by protein phosphatase 2B (calcineurin). *J Biol Chem.* 1999; 274(21):15173-15180. (Biology)

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	800.979.9408	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD

