



43-2040 5 mg
 Biotinylated
 Peroxidase
 Lot No.

BIOTINYLATED PEROXIDASE

BIOTIN/AVIDIN/STREPTAVIDIN REAGENTS

FORM:

2 ml liquid. This product is in 10 mM phosphate buffered saline, pH 7.4 containing 1% bovine serum albumin and the protein concentration of biotinylated peroxidase is 2.5 mg/ml. Purified horseradish peroxidase (RZ>3) is biotinylated under optimum conditions and is free of unconjugated biotin.

RECOMMENDED USAGE FOR EIA:

Biotinylated peroxidase (1 mg/ml)	5 µl
Streptavidin (1 mg/ml)	10 µl
PBS,(10 mM, pH 7.4) containing 0.05% Tween 20	25 ml

Mix well and incubate for 30 minutes at room temperature before use. Discard unused diluted complex after 24 hours. Note: For histochemical staining, use 5 to 10 times more concentrated complex.

STORAGE:

Store at 4°C.

REMARKS: For research use only. Not for human or drug use. Product contains 0.1% proclin as preservative.

www.invitrogen.com

Invitrogen Corporation • 542 Flynn Rd • Camarillo • CA 93012 • Tel: 800.955.6288
 E-mail: techsupport@invitrogen.com

PI 43-2040

(Rev 10/08) DCC-08-1089

Important Licensing Information - These products may be covered by one or more Limited Use Label Licenses (see the Invitrogen Catalog or our website, www.invitrogen.com). By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.



43-2040 5 mg
 Biotinylated
 Peroxidase
 Lot No.

BIOTINYLATED PEROXIDASE

BIOTIN/AVIDIN/STREPTAVIDIN REAGENTS

FORM:

2 ml liquid. This product is in 10 mM phosphate buffered saline, pH 7.4 containing 1% bovine serum albumin and the protein concentration of biotinylated peroxidase is 2.5 mg/ml. Purified horseradish peroxidase (RZ>3) is biotinylated under optimum conditions and is free of unconjugated biotin.

RECOMMENDED USAGE FOR EIA:

Biotinylated peroxidase (1 mg/ml)	5 µl
Streptavidin (1 mg/ml)	10 µl
PBS,(10 mM, pH 7.4) containing 0.05% Tween 20	25 ml

Mix well and incubate for 30 minutes at room temperature before use. Discard unused diluted complex after 24 hours. Note: For histochemical staining, use 5 to 10 times more concentrated complex.

STORAGE:

Store at 4°C.

REMARKS: For research use only. Not for human or drug use. Product contains 0.1% proclin as preservative.

www.invitrogen.com

Invitrogen Corporation • 542 Flynn Rd • Camarillo • CA 93012 • Tel: 800.955.6288
 E-mail: techsupport@invitrogen.com

PI 43-2040

(Rev 10/08) DCC-08-1089

Important Licensing Information - These products may be covered by one or more Limited Use Label Licenses (see the Invitrogen Catalog or our website, www.invitrogen.com). By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.