

Technical Data Sheet


APC Mouse Anti-Human CD196 (CCR6)

Product Information

Material Number:	560619
Alternate Name:	BN-1; C-C chemokine receptor type 6; C-C CKR-6; CC-CKR-6; CCR-6; CD196
Size:	50 tests
Vol. per Test:	5 µl
Clone:	11A9
Isotype:	Mouse IgG1, κ
Reactivity:	QC Testing: Human
Workshop:	HLDA VII
Storage Buffer:	Aqueous buffered solution containing BSA and ≤0.09% sodium azide.

Description

CCR6 is a seven-transmembrane, G-protein-coupled, glycoprotein receptor that is a member of the beta chemokine receptor family. CCR6 was clustered as CD196 in the 7th HLDA Workshop. The human CCR6 gene has been mapped to chromosome 6q27. CCR6 is a receptor for the CC chemokine CCL20/MIP-3α/LARC/Exodus and also binds with lower affinity to and mediates responses to beta-defensin2/hBD-2. CCR6 is predominantly expressed by B lymphocytes, certain subsets of effector and memory T cells and by immature dendritic cells but not by monocytes, NK cells, or granulocytes. Skin-homing CLA (Cutaneous Lymphocyte Antigen)-positive memory T cells, Th1 cells, regulatory T cells and IL-17A-producing Th17 cells predominantly express high levels of CCR6. CCR6 mediates the trafficking of T, B, and dendritic cells to epithelial sites near the skin and mucosal surfaces during inflammatory and immunological responses. The monoclonal antibody 11A9 reacts with the human CC chemokine receptor, CCR6. An N-terminal peptide of human CCR6 was used as an immunogen to generate the 11A9 hybridoma. The 11A9 antibody does not cross-react with human CCR1, CCR2, CCR3, CCR4, CCR5, CCR7, CCR8, CCR9, CXCR1, CXCR2, CXCR3, CXCR4 and CXCR5 receptors. This antibody is NOT a neutralizing antibody.



Flow cytometric analysis of CD196 on lysed whole blood. Human lysed whole blood was stained with the APC Mouse Anti-Human CD196 antibody (unshaded) or with a APC Mouse IgG1, κ isotype control (shaded). Histograms were derived from gated events based on light scattering characteristics for lymphocytes. Flow cytometry was performed on a BD™ LSR II flow cytometry system.

Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated to APC under optimum conditions, and unconjugated antibody and free APC were removed.

Application Notes

Application

Flow cytometry	Routinely Tested
----------------	------------------

Recommended Assay Procedure:

Flow cytometry: Chemokine receptors are known to internalize during manipulation resulting in low frequency expression. Investigators are advised to perform immunophenotyping studies of chemokine receptors on freshly collected samples (<24 Hrs). Incubation with the antibody should be done at 4°C in the dark. Cellular manipulation, such as Ficoll separation, freezing, or exposure to cold temperatures prior to staining should be minimized and have been shown to cause a decrease in staining intensity and/or inconsistent results.

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2008 BD



Investigators should note that alternative staining procedures may be necessary. A multiple-step staining procedure may be helpful, in some instances, to amplify immunofluorescent signals for the flow cytometric analysis of CD196 expression. Investigators may find the Purified Mouse Anti-Human CD196 antibody (MN 559560) to be useful in conjunction with appropriate secondary and tertiary reagents for detecting low frequency expression, such as with Biotin Goat Anti-Mouse Ig (MN 553999) and PE Streptavidin (MN 554061) or APC Streptavidin (MN 554067).

Suggested Companion Products

Catalog Number	Name	Size	Clone
555751	APC Mouse IgG1, κ Isotype Control	100 tests	MOPC-21
555899	Lysing Buffer	100 ml	(none)
559560	Purified Mouse Anti-Human CD196 (CCR6)	0.5 mg	11A9
553999	Biotin Goat Anti-Mouse Ig (Multiple Adsorption)	0.5 mg	Polyclonal
554061	PE Streptavidin	0.5 mg	(none)
554067	APC Streptavidin	0.1 mg	(none)

Product Notices

1. This reagent has been pre-diluted for use at the recommended Volume per Test. We typically use 1×10^6 cells in a 100- μ l experimental sample (a test).
2. An isotype control should be used at the same concentration as the antibody of interest.
3. This APC-conjugated reagent can be used in any flow cytometer equipped with a dye, HeNe, or red diode laser.
4. Source of all serum proteins is from USDA inspected abattoirs located in the United States.
5. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
6. For fluorochrome spectra and suitable instrument settings, please refer to our Fluorochrome Web Page at www.bdbiosciences.com/colors.
7. Please refer to www.bdbiosciences.com/pharming/protocols for technical protocols.

References

Baba M, Imai T, Nishimura M, et al. Identification of CCR6, the specific receptor for a novel lymphocyte-directed CC chemokine LARC. *J Biol Chem.* 1997; 272(23):14893-14898. (Biology)

Greaves DR, Wang W, Dairaghi DJ, et al. CCR6, a CC chemokine receptor that interacts with macrophage inflammatory protein 3 α and is highly expressed in human dendritic cells. *J Exp Med.* 1997; 186(6):837-844. (Biology)

Homey B, Dieu-Nosjean MC, Wiesenborn A, et al.. Up-regulation of macrophage inflammatory protein-3 α /CCL20 and CC chemokine receptor 6 in psoriasis.. *J Immunol.* 2000; 164(12):6621-6632. (Biology)

Kim CH, Rott L, Kunkel EJ, et al.. Rules of chemokine receptor association with T cell polarization in vivo.. *J Clin Invest.* 2001; 108(9):1331-1339. (Biology)

Liao F, Alderson R, Su J, Ullrich SJ, Kreider BL, Farber JM. STRL22 is a receptor for the CC chemokine MIP-3 α . *Biochem Biophys Res Commun.* 1997; 236(1):212-217. (Biology)

Liao F, Lee HH, Farber JM. Cloning of STRL22, a new human gene encoding a G-protein-coupled receptor related to chemokine receptors and located on chromosome 6q27. *Genomics.* 1997; 40(1):175-180. (Biology)

Liao F, Rabin RL, Smith CS, Sharma G, Nutman TB, Farber JM. CC-chemokine receptor 6 is expressed on diverse memory subsets of T cells and determines responsiveness to macrophage inflammatory protein 3 α . *J Immunol.* 1999; 162(1):186-194. (Biology)

Lim HW, Lee J, Hillsamer P, Kim CH.. Human Th17 cells share major trafficking receptors with both polarized effector T cells and FOXP3+ regulatory T cells.. *J Immunol.* 2008; 180(1):122-129. (Biology)

Power CA, Church DJ, Meyer A, et al. Cloning and characterization of a specific receptor for the novel CC chemokine MIP-3 α from lung dendritic cells. *J Exp Med.* 1997; 186(6):825-835. (Biology)

Yang D, Chertov O, Bykovskaia SN, et al.. Beta-defensins: linking innate and adaptive immunity through dendritic and T cell CCR6.. *Science.* 1999; 286(5439):525-528. (Biology)

Zaballos A, Varona R, Gutierrez J, Lind P, Marquez G. Molecular cloning and RNA expression of two new human chemokine receptor-like genes. *Biochem Biophys Res Commun.* 1996; 227(3):846-853. (Biology)