

Technical Data Sheet



APC Rat Anti-Mouse Vα2 TCR

Product Information

Material Number:	560622
Size:	50 µg
Concentration:	0.2 mg/ml
Clone:	B20.1
Immunogen:	Soluble αβ TCR from mouse cytotoxic T-cell clone KB5-C20
Isotype:	Rat (LOU) IgG2a, λ
Reactivity:	QC Testing: Mouse
Storage Buffer:	Aqueous buffered solution containing ≤0.09% sodium azide.

Description

The B20.1 monoclonal antibody specifically binds to most members of the Vα2 T-cell Receptor (TCR) subfamily in mice having the *a*, *b*, and *c* haplotypes of the *Tcrb* gene complex. B20.1 antibody may crossreact with Vδ8 TCR, which shares >90% sequence homology with Vα2 TCR. Levels of B20.1+ T cells appear to be influenced by Vα haplotypes. Moreover, the frequencies of Vα2+ CD8+ and CD4+ T cells are influenced by H-2 haplotypes.



Flow cytometric analysis of Vα2 TcR on mouse lymph node cells. Lymph node cells from BALB/c mice were stained with FITC Rat Anti-Mouse CD4 (Cat. No. 553046) and FITC Rat Anti-Mouse CD8b.2 (Cat. No. 553040) in addition to either a APC Rat IgG2a, λ isotype control (left panel) or with the APC Rat Anti-Mouse Vα2 TcR antibody (right panel). Dot plots were derived from gated events based on light scattering characteristics for lymph node cells. Flow cytometry was performed on a BD™ LSR II flow cytometry system.

Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated to APC under optimum conditions, and unconjugated antibody and free APC were removed.

Application Notes

Application

Flow cytometry	Routinely Tested
----------------	------------------

Suggested Companion Products

Catalog Number	Name	Size	Clone
560720	APC Rat IgG2a, λ Isotype Control	0.1 mg	B39-4
553141	Purified Rat Anti-Mouse CD16/CD32 (Mouse BD Fc Block™)	0.1 mg	2.4G2
553046	FITC Rat Anti-Mouse CD4	0.1 mg	RM4-5
553040	FITC Rat Anti-Mouse CD8b.2	0.5 mg	53-5.8

Product Notices

- Since applications vary, each investigator should titrate the reagent to obtain optimal results.
- An isotype control should be used at the same concentration as the antibody of interest.
- This APC-conjugated reagent can be used in any flow cytometer equipped with a dye, HeNe, or red diode laser.
- Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

BD Biosciences

bdbiosciences.com

United States 877.232.8995 Canada 888.268.5430 Europe 32.53.720.550 Japan 0120.8555.90 Asia Pacific 65.6861.0633 Latin America/Caribbean 0800.771.7157

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2011 BD



5. For fluorochrome spectra and suitable instrument settings, please refer to our Fluorochrome Web Page at www.bdbiosciences.com/colors.
6. Please refer to www.bdbiosciences.com/pharming/protocols for technical protocols.

References

- Gregoire C, Rebai N, Schweisguth F, et al. Engineered secreted T-cell receptor alpha beta heterodimers. *Proc Natl Acad Sci U S A*. 1991; 88(18):8077-8081. (Biology)
- Pircher H, Rebai N, Groettrup M, et al. Preferential positive selection of V alpha 2+ CD8+ T cells in mouse strains expressing both H-2k and T cell receptor V alpha a haplotypes: determination with a V alpha 2-specific monoclonal antibody. *Eur J Immunol*. 1992; 22(2):399-404. (Immunogen)
- Tomonari K, Fairchild S, Rosenwasser OA. Influence of viral superantigens on V beta- and V alpha-specific positive and negative selection. *Immunol Rev*. 1993; 131:131-168. (Biology)