

## Technical Data Sheet


## FITC Rat Anti-Mouse CD335 (NKp46)

## Product Information

<b>Material Number:</b>	560756
<b>Alternate Name:</b>	NKp46; Ar1; Ly94; Lymphocyte antigen 94; Mar1; MAR-1; Mouse activating rece
<b>Size:</b>	0.1 mg
<b>Concentration:</b>	0.5 mg/ml
<b>Clone:</b>	29A1.4
<b>Isotype:</b>	Rat IgG2a, $\kappa$
<b>Reactivity:</b>	QC Testing: Mouse
<b>Storage Buffer:</b>	Aqueous buffered solution containing $\leq 0.09\%$ sodium azide.

## Description

The monoclonal antibody 29A1.4 specifically binds to mouse CD335, also known as NKp46. NKp46 is a 46 kDa type I transmembrane glycoprotein that is a member of the natural cytotoxicity receptor (NCR) family and immunoglobulin superfamily. NKp46 is encoded by the Ncr1 gene located on chromosome 7. NKp46 functions as a cytotoxicity triggering receptor and is selectively expressed by immature and mature NK cells in all mouse strains tested. NKp46 is detected on a minute fraction of NK-like T cells (less than 2% of NKp46<sup>+</sup> express CD3e) but not on CD1d-restricted NKT cells from C57BL/6 mice. When immobilized on tissue culture plates, the 29A1.4 antibody reportedly stimulates NK cells to produce interferon-gamma and to release their cytoplasmic granule contents. Although the ligands for the NKp46 receptor have not been fully characterized, recent evidence indicates that this receptor plays an important role in the NK cell-mediated recognition and killing of some virus-infected cells and tumor cells. The immunogen used to generate the 29A1.4 clone was mouse NKp46-Fc recombinant protein.



**Flow cytometric analysis of FITC anti-mouse CD335 (NKp46) expression on mouse splenocytes.** C57BL/6 and BALB/c mouse spleen cells were stained separately with FITC anti-mouse CD335 (NKp46) antibody. After washing, C57BL/6 cells were stained with PE anti-mouse NK-1.1 (NKR-P1B and NKR-P1C) antibody (Cat. No. 553165; left panel) and BALB/c cells were stained with PE anti-mouse CD49b (DX5) (Cat. No. 553858; right panel). Two-color dot plots showing the correlated expression patterns of CD335/NKp46 and either NK1.1/CD161 (C57BL/6 cells; left panel) or DX5/CD49b (BALB/c cells; right panel) were derived from gated events with the forward and side light-scatter characteristics of viable lymphocytes. Flow cytometry was performed using a BD™ LSRII System.

## Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with FITC under optimum conditions, and unreacted FITC was removed.

## BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.268.5430	32.53.720.550	0120.8555.90	65.6861.0633	0800.771.7157

For country-specific contact information, visit [bdbiosciences.com/how\\_to\\_order/](http://bdbiosciences.com/how_to_order/)

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2011 BD



## Application Notes

### Application

Flow cytometry

Routinely Tested

### Suggested Companion Products

Catalog Number	Name	Size	Clone
553165	PE Mouse Anti-Mouse NK-1.1	0.2 mg	PK136
553858	PE Rat Anti-Mouse CD49b	0.2 mg	DX5

### Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
3. For fluorochrome spectra and suitable instrument settings, please refer to our Fluorochrome Web Page at [www.bdbiosciences.com/colors](http://www.bdbiosciences.com/colors).
4. An isotype control should be used at the same concentration as the antibody of interest.
5. Please refer to [www.bdbiosciences.com/pharming/en/protocols](http://www.bdbiosciences.com/pharming/en/protocols) for technical protocols.

### References

Biassoni R, Pessino A, Bottino C, Pende D, Moretta L, Moretta A. The murine homologue of the human NKp46, a triggering receptor involved in the induction of natural cytotoxicity. *Eur J Immunol.* 1999; 29(3):1014-1020. (Biology)

Gazit R, Gruda R, Elboim M, et al. Lethal influenza infection in the absence of the natural killer cell receptor gene Ncr1. *Nat Immunol.* 2006; 7(5):517-523. (Biology)

Joncker NT, Fernandez NC, Treiner E, Vivier E, Raulet DH. NK cell responsiveness is tuned commensurate with the number of inhibitory receptors for self-MHC class I: the rheostat model. *J Immunol.* 2009; 182(8):4572-4580. (Clone-specific: Flow cytometry)

Walzer T, Blery M, Chaix J, et al. Identification, activation, and selective in vivo ablation of mouse NK cells via NKp46. *Proc Natl Acad Sci U S A.* 2007; 104(9):3384-3389. (Clone-specific: Activation, Flow cytometry)