## Technical Data Sheet

# PE Mouse Anti-Human IL-9

## **Product Information**

Material Number:	560814
Size:	25 tests
Vol. per Test:	5ul
Clone:	MH9A3
Immunogen:	Human IL-9 Recombinant Protein
Isotype:	Mouse (C57BL/6) IgG1, ĸ
Reactivity:	Reported: Human
Storage Buffer:	Aqueous buffered solution containing BSA and ≤0.09% sodium azide.

## Description

The MH9A3 monoclonal antibody specifically binds to human interleukin-9 (IL-9). Human IL-9 is a multifunctional cytokine and a member of the type I cytokine (hematopoietin) family that includes IL-2, IL-4, IL-7, IL-15 and IL-21. This cytokine is encoded by the *IL9* gene that is resident on chromosome 5q31.1. IL-9 is expressed by activated CD4-positive T helper cells, by some transformed T cells and by eosinophils, mast cells and neutrophils. IL-9 induces the proliferation, differentiation, and effector function of various cell types including T lymphocytes, B lymphocytes, mast cells, eosinophils, neutrophils, hematopoietic cells and epithelial cells. It potentiates the interleukin-4-induced IgM, IgG and IgE responses by human B lymphocytes. IL-9 has been implicated in human allergic disorders such as asthma and malignancies such as Hodgkin's disease. IL-9 exerts its biological activities through binding to the surface IL-9 receptor (IL-9R) complex comprised of the IL-9R alpha subunit (IL-9R $\alpha$ ; CD129) and the common cytokine receptor gamma subunit ( $\gamma$ ; CD132). IL-9 signaling through its receptor includes activation of the Janus kinases 1 and 3 (JAK1 and JAK3) and activation of Signal transducer and activator of transcription 1, 3 and 5 factors (STAT1, STAT3 and STAT5).



Flow cytometric analysis of IL-9 in stimulated human CD4-positive T cells. Human peripheral blood mononuclear cells were stimulated in a tissue culture plate coated with NA/LE Mouse Anti-Human CD3 (Cat. No. 555329; 10 µg/ml, coated overnight at 4°C) and soluble NA/LE Mouse Anti-Human CD28 (Cat. No. 555725; 1 µg/ml) antibodies plus recombinant Human IL-2 (Cat. No. 554603; 10 ng/ml), IL-4 (Cat. No. 554605; 50 ng/ml), and TGF-β (Cat. No. 356039; 10 ng/ml) proteins and NA/LE Mouse Anti-Human IFN-γ (Cat. No. 554698; 10 µg/ml) antibody for 5 days. The cells were harvested and restimulated with PMA (Sigma P8139; 50 ng/ml) and ionomycin (Sigma I9657; 1 µg/ml) in the presence of BD GolgiStop™ Protein Transport Inhibitor (Cat. No. 554724) for 5 hours. The cells were then fixed and permeabilized using the BD Cytofix/Cytoperm™ Fixation/Permeablization Kit (Cat. No. 554714) followed by staining with PE Mouse Anti-Human IL-9 (Cat. No. 560814) and APC Mouse Anti-Human CD4 (Cat. No. 555349) antibodies. Two-color flow cytometric dot plots showing correlated expression patterns of CD4 versus IL-9 were derived from CD4-positive cell gated events with the forward and side light-scatter characteristics of viable lymphocytes. Flow cytometry was performed on a BD LSRII™ System.

#### **BD Biosciences**

bdbiosciences.com United States Canada Asia Pacific Latin America/Caribbean Europe Japan 32.53.720.550 0120.8555.90 877.232.8995 888.268.5430 65.6861.0633 0800.771.7157 For country-specific contact information, visit bdbiosciences.com/how\_to\_order/ Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited. For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale. BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2011 BD



## **Preparation and Storage**

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. The antibody was conjugated with R-PE under optimum conditions, and unconjugated antibody and free PE were removed. Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

## **Application Notes**

Application					
Intracellular staining (	flow cytometry) Routinely Tested	Routinely Tested			
Suggested Companion Products					
Catalog Number	Name	Size	Clone		
551436	PE Mouse IgG1 Kappa Isotype Control	50 tests	MOPC-21		
555349	APC Mouse Anti-Human CD4	100 tests	RPA-T4		
554722	Fixation and Permeabilization Solution	125 ml	(none)		
554723	Perm/Wash Buffer	100 ml	(none)		
554655	Fixation Buffer	100 ml	(none)		
555029	Protein Transport Inhibitor (Containing Brefeldin A)	1.0 ml	(none)		
554724	Protein Transport Inhibitor (Containing Monensin)	0.7 ml	(none)		
554656	Stain Buffer (FBS)	500 ml	(none)		
554657	Stain Buffer (BSA)	500 ml	(none)		
554714	BD Cytofix/Cytoperm <sup>™</sup> Fixation/Permeablization Kit	250 tests	(none)		
554603	Recombinant Human IL-2	10 µg	(none)		
554605	Recombinant Human IL-4	5 µg	(none)		

555329	Purified NA/LE Mouse Anti-Human CD3	0.5 mg	UCHT1
555725	Purified NA/LE Mouse Anti-Human CD28	0.5 mg	CD28.2
554698	Purified NA/LE Mouse Anti-Human IFN-y	0.5 mg	B27
354039	Transforming Growth Factor-beta (TGF-b), human natural, 1 µg	10 tests	hTGF- beta
356039	Transforming Growth Factor-b (TGF-b), human natural, 1 X 5 $\mu$ g	NA	(none)
356040	Transforming Growth Factor-b (TGF-b), human natural, 5 X 2 $\mu$ g	NA	(none)

#### **Product Notices**

1. This reagent has been pre-diluted for use at the recommended Volume per Test. We typically use  $1 \times 10^{6}$  cells in a 100-µl experimental sample (a test).

- 2. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
- 3. For fluorochrome spectra and suitable instrument settings, please refer to our Fluorochrome Web Page at www.bdbiosciences.com/colors.
- 4. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
- 5. Source of all serum proteins is from USDA inspected abattoirs located in the United States.

### References

Demoulin JB, Van Roost E, Stevens M, Groner B, Renauld JC. Distinct roles for STAT1, STAT3, and STAT5 in differentiation gene induction and apoptosis inhibition by interleukin-9. *J Biol Chem.* 1999; 274:25855-25861. (Biology)

Dugas B, Renauld JC, Pène J, Bonnefoy JY, Peti-Frère C, Braquet P, Bousquet J, Van Snick J, Mencia-Huerta JM. Interleukin-9 potentiates the interleukin-4-induced immunoglobulin (IgG, IgM and IgE) production by normal human B lymphocytes. *Eur J Immunol.* 1993 July; 23(7):1687-1692. (Biology) Houssiau FA, Schandene L, Stevens M. A cascade of cytokines is responsible for IL-9 expression in human T cells. Involvement of IL-2, IL-4, and IL-10. *J Immunol.* 1995; 154(6):2624-2630. (Biology)

Jenmalm MC, Van Snick J, Cormont F, Salman B. Allergen-induced Th1 and Th2 cytokine secretion in relation to specific allergen sensitization and atopic symptoms in children. *Clin Exp Allergy*. 2001 October; 31(10):1528-1535. (Clone-specific: ELISA)

Knoops L, Renauld JC. IL-9 and its receptor: from signal transduction to tumorigenesis. Growth Factors. 2004; 22:207-215. (Biology)

Merz H, Houssiau FA, Orscheschek K, Renauld JC, Fliedner A, Herin M, Noel H, Kadin M, Mueller-Hermelink HK, Van Snick J, et al. Interleukin-9 expression in human malignant lymphomas: unique association with Hodgkin's disease and large cell anaplastic lymphoma. *Blood.* 1991 September; 78(5):1311-1317. (Biology)

Renauld JC. New insights into the role of cytokines in asthma. J Clin Pathol. 2001 August; 54(8):577-589. (Biology)

Soler D, Chapman TR, Poisson LR. CCR8 expression identifies CD4 memory T cells enriched for FOXP3+ regulatory and Th2 effector lymphocytes. J Immunol. 2006; 177(10):6940-6951. (Biology)

Soroosh P, Doherty TA. Th9 and allergic disease. Immunology. 2009; 127(4):450-458. (Biology)