Technical Data Sheet


PerCP-Cy™5.5 Mouse Anti-Human CD69

Product Information

Material Number:	560738
Alternate Name:	Very Early Activation Antigen
Size:	50 tests
Vol. per Test:	5 μl
Clone:	FN50
Isotype:	Mouse IgG1, ĸ
Reactivity:	Human
	QC Testing: Rhesus or Baboon or Cynomolgus
Workshop:	IV A091
Storage Buffer:	Aqueous buffered solution containing BSA and ≤0.09% sodium azide.

Description

Reacts with a 28/34 kDa dimeric glycoprotein expressed early during activation of lymphocytes and monocytes. FN50 monoclonal antibody labels NK cells and most lymphocytes of the follicular mantle and perifollicular/interfollicular zone as well as intragerminal center T cells of lymph nodes and tonsils.



Flow cytometric analysis for CD69 in stimulated Rhesus macaque peripheral blood mononuclear cells (PBMC). PBMC from Rhesus macaque were stimulated for at least 4-6 hours with 20 ng/mL PMA (Sigma-Aldrich Cat. No. P-8139) and 250 ng/mL calcium ionophore A23187 (Sigma-Aldrich Cat. No. C-9275). Cells were then stained with either a PerCP-CyTM5.5 Mouse IgG1, κ isotype control (shaded) or with the PerCP-CyTM5.5 Mouse Anti-Human CD69 antibody (unshaded). Histograms were derived from gated events based on light scattering characteristics for Jymphocytes. Flow cytometry was performed on a BDTM LSR II flow cytometry system.

500 ml

(none)

Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with PerCP-Cy5.5 under optimum conditions, and unconjugated antibody and free PerCP-Cy5.5 were removed. Storage of PerCP-Cy5.5 conjugates in unoptimized diluent is not recommended and may result in loss of signal intensity.

Application Notes

Application				
Flow cytometry	Routinely Tested			
Suggested Compan	ion Products			
Catalog Number	Name	Size	Clone	
555899	Lysing Buffer	100 ml	(none)	
550795	PerCP-Cy [™] 5.5 Mouse IgG1 κ Isotype Control	0.1 mg	MOPC-21	

Product Notices

554656

- 1. This reagent has been pre-diluted for use at the recommended Volume per Test. We typically use 1×10^{6} cells in a 100-µl experimental sample (a test).
- 2. An isotype control should be used at the same concentration as the antibody of interest.

Stain Buffer (FBS)

3. Please observe the following precautions: Absorption of visible light can significantly alter the energy transfer occurring in any tandem fluorochrome conjugate; therefore, we recommend that special precautions be taken (such as wrapping vials, tubes, or racks in aluminum foil) to prevent exposure of conjugated reagents, including cells stained with those reagents, to room illumination.

BD Biosciences

bdbiosciences.com							
United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean		
877.232.8995	800.979.9408	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995		
For country co	ntact informatio	on, visit bdbiosci	ences.com/conta	ct			
of any patents. BL	D Biosciences will n	ot be help responsi	ble for patent infrin	gement or other vio	ise the above product in violatio		
use of our product	ts. Purchase does n	not include or carry	any right to resell ou	r transfer this produ	lations that may occur with the		
product or as a co	imponent of anoth	er product. Any use	of this product oth	er than the permitte	ct either as a stand-alone		
written authoriza	tion of Becton, Dic	kinson and Compar	ny is stictly prohibite	d.	id use without the express		
For Research Use	Only. Not for use in	n diagnostic or there	apeutic procedures.	Not for resale.	n and Company. © 2011 BD		

- Cy is a trademark of Amersham Biosciences Limited. This conjugated product is sold under license to the following patents: US Patent Nos. 4. 5,486,616; 5,569,587; 5,569,766; 5,627,027.
- This product is subject to proprietary rights of Amersham Biosciences Corp. and Carnegie Mellon University and made and sold under 5. license from Amersham Biosciences Corp. This product is licensed for sale only for research. It is not licensed for any other use. If you require a commercial license to use this product and do not have one return this material, unopened to BD Biosciences, 10975 Torreyana Rd, San Diego, CA 92121 and any money paid for the material will be refunded.
- 6. PerCP-Cy5.5 is optimized for use with a single argon ion laser emitting 488-nm light. Because of the broad absorption spectrum of the tandem fluorochrome, extra care must be taken when using dual-laser cytometers, which may directly excite both PerCP and Cy5.5TM. We recommend the use of cross-beam compensation during data acquisition or software compensation during data analysis.
- 7. PerCP-Cy5.5-labelled antibodies can be used with FITC- and R-PE-labelled reagents in single-laser flow cytometers with no significant spectral overlap of PerCP-Cy5.5, FITC, and R-PE fluorescence.
- 8. Source of all serum proteins is from USDA inspected abattoirs located in the United States.
- 9. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
- For fluorochrome spectra and suitable instrument settings, please refer to our Fluorochrome Web Page at www.bdbiosciences.com/colors. 10.
- Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols. 11.

References

Knapp W, Dorken B, Rieber EP, et al, ed. Leucocyte Typing IV. New York: Oxford University Press; 1989:1-1208. (Biology) Schlossman S, Boumell L, et al, ed. Leucocyte Typing V. New York: Oxford University Press; 1995. (Biology)

BD Biosciences

bdbiosciences.com

United States Canada Europe 877.232.8995 800.979.9408 32.53.720.550 0120.8555.90

65.6861.0633 For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be constructed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be help responsible for patent infringement or other violations that may occur with the use of our products. Buckness will not be help responsible for patent infringement or other violations that may occur with the product or as a component of another product. Any use of this product other than the permitted use without the express

Asia Pacific

Latin America/Caribbean

55.11.5185.9995

written authorization of Becton, Dickinson and Company is stictly prohibited. For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale. Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD

Japan

