## **Technical Data Sheet**

# PE Mouse Anti-Human CD11b/Mac-1

#### **Product Information**

**Material Number:** 

Mac-1α, integrin αM subunit, CR3 α chain Alternate Name:

25 tests Size: 20 ul Vol. per Test:

ICRF44 (also known as 44) Clone:


Mouse IgG1, κ Isotype: Reactivity: QC Testing: Human

IV M047 Workshop:


Aqueous buffered solution containing BSA and ≤0.09% sodium azide. Storage Buffer:

#### Description

The ICRF44 monoclonal antibody reacts with CD11b, the 165-kDa adhesion glycoprotein that associates with the 95-kDa integrin β2 (CD18) to form the CD11b/CD18 complex, also known as Mac-1 or CR3. CD11b is expressed on activated lymphocytes, monocytes, granulocytes, and a subset of NK cells. CD11b functions in cell-cell and cell-substrate interactions and is a receptor for iC3b, CD54 (ICAM-1), CD102 (ICAM-2) and CD50 (ICAM-3). This antibody significantly inhibits polymorphonuclear leukocyte aggregation in response to fMLP.



Profile of lymphocytes analyzed on a FACScan (BDIS, San Jose, CA)



Profile of granulocytes analyzed on a FACScan (BDIS, San Jose, CA)

### **Preparation and Storage**

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with R-PE under optimum conditions, and unconjugated antibody and free PE were removed. Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

### **Application Notes**

### Application

Flow cytometry Routinely Tested

### **Suggested Companion Products**

<u>Clone</u> 555749 PE Mouse IgG1, κ Isotype Control 100 tests MOPC-21

### **Product Notices**

- This reagent has been pre-diluted for use at the recommended Volume per Test. We typically use  $1 \times 10^6$  cells in a 100- $\mu$ l experimental
- Since applications vary, each investigator should titrate the reagent to obtain optimal results.

## **BD Biosciences**

bdbiosciences.com

**United States** 877.232.8995 888.268.5430 32.53.720.550 0120.8555.90 65.6861.0633 0800.771.7157

For country-specific contact information, visit bdbiosciences.com/how\_to\_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.
For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.
BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2011 BD



- 3. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
- 4. For fluorochrome spectra and suitable instrument settings, please refer to our Fluorochrome Web Page at www.bdbiosciences.com/colors.
- 5. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
- 6. Source of all serum proteins is from USDA inspected abattoirs located in the United States.

#### References

Barclay NA, Brown MH, Birkeland ML, et al, ed. *The Leukocyte Antigen FactsBook*. San Diego, CA: Academic Press; 1997. (Biology) Knapp W, Dorken B, Rieber EP, et al, ed. *Leucocyte Typing IV*. New York: Oxford University Press; 1989:1-1208. (Clone-specific)

561001 Rev. 1 Page 2 of 2