

## Technical Data Sheet


## PE Mouse anti-Human IL-17F

## Product Information

<b>Material Number:</b>	<b>561198</b>
<b>Alternate Name:</b>	Interleukin-17F; cytokine ML-1; ML-1; IL-24
<b>Size:</b>	25 tests
<b>Vol. per Test:</b>	5 µl
<b>Clone:</b>	O33-782
<b>Immunogen:</b>	Human IL-17F
<b>Isotype:</b>	Mouse IgG1, κ
<b>Reactivity:</b>	QC Testing: Human
<b>Storage Buffer:</b>	Aqueous buffered solution containing BSA and ≤0.09% sodium azide.

## Description

The O33-782 monoclonal antibody specifically binds to Interleukin-17F (IL-17F). IL-17F is a member of the IL-17 family of cytokines. IL-17F is encoded by the *IL17F* gene located in chromosome 6 (location: 6p12). IL-17F is a proinflammatory cytokine that is produced by activated T cells including differentiated CD4+ T helper 17 (Th17) cells. Activated Th17 cells can express disulfide-linked IL-17F and IL-17A homodimers as well as IL-17A/IL-17F heterodimers. These IL-17 dimers act by binding to and signaling through IL-17 receptor complexes (IL-17R). IL-17R are comprised of transmembrane IL-17RA and IL-17RC protein subunits that are expressed by a variety of target cells including epithelial and endothelial cells, keratinocytes, fibroblasts, and granulocytes. IL-17F can induce target cells to produce proinflammatory cytokines such as IL-1β, IL-6, G-CSF, GM-CSF, and TNF and chemokines including CXCL1/Gro-α, CXCL2/Gro-β, and CXCL8/IL-8 that attract and activate leukocytes, eg, neutrophils. Th17 and other IL-17F-producing cells play protective roles in the clearance of extracellular pathogens, including bacteria and fungi. IL-17F can also play adverse roles in inflammation associated with asthma and autoimmune diseases.



**Flow cytometric analysis of IL-17F expression by resting and activated human peripheral blood CD4+ T cells and Th17 polarized cells.** Human peripheral blood mononuclear cells were either unstimulated (Left Panels) or stimulated with Phorbol 12-Myristate 13-Acetate (PMA; Sigma P-8139) plus Ionomycin (Sigma; I-0634) in the presence of BD GolgiStop™ Protein Transport Inhibitor (Cat. No. 554724) for 5 hours (Middle Panels) or were cultured in Th17 polarization conditions and restimulated with PMA and Ionomycin in the presence of BD GolgiStop™ for 5 hours (Right Panels). Cells were then fixed and permeabilized using BD Cytofix/Cytoperm™ reagents (Cat. No. 554714) followed by staining with PE Mouse anti-Human IL-17F (Cat. No. 561198), PerCP-Cy5.5 Mouse anti-Human CD4 (Cat. No. 341654), and FITC Mouse Anti-Human IFN-γ (Cat. No. 554700) or Alexa Fluor® 647 Mouse anti-Human IL-17A (Cat. No. 560490). Two-color flow cytometric dot plots showing the correlated expression patterns of IL-17F versus IL-17A or IFN-γ were derived from CD4+ gated events with the forward and side light-scatter characteristics of intact lymphocytes. Flow cytometry was performed using a BD™ LSR II Flow Cytometer System. Other compatible fixation and permeabilization treatments are listed in the "Recommended Assay Procedure."

## BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.268.5430	32.53.720.550	0120.8555.90	65.6861.0633	0800.771.7157

For country-specific contact information, visit [bdbiosciences.com/how\\_to\\_order/](http://bdbiosciences.com/how_to_order/)

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2011 BD



## Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with R-PE under optimum conditions, and unconjugated antibody and free PE were removed.

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

## Application Notes

### Application

Intracellular staining (flow cytometry)	Routinely Tested
-----------------------------------------	------------------

### Recommended Assay Procedure:

This antibody conjugate is suitable for intracellular staining of human peripheral blood mononuclear cells using BD Cytotfix/Cytoperm™ reagents, BD Pharmingen™ Human FoxP3 Buffer Set or BD™ Phosflow fixation and permeabilization buffers (Fix buffer I with Perm/Wash Buffer I, Perm Buffer II, or Perm Buffer III).

### Suggested Companion Products

Catalog Number	Name	Size	Clone
554724	Protein Transport Inhibitor (Containing Monensin)	0.7 ml	(none)
554714	BD Cytotfix/Cytoperm™ Fixation/Permeabilization Kit	250 tests	(none)
341654	PerCP-Cy5.5 Mouse anti-Human CD4	50 tests	SK3
554700	FITC Mouse Anti-Human IFN-γ	0.1 mg	B27
560490	Alexa Fluor® 647 Mouse anti-Human IL-17A	100 tests	N49-653
554680	PE Mouse IgG1, κ Isotype Control	0.1 mg	MOPC-21
557870	Fix Buffer I	250 ml	(none)
557885	Perm/Wash Buffer I	125 ml	(none)
558052	Perm Buffer II	125 ml	(none)
558050	Perm Buffer III	125 ml	(none)
560098	Human FoxP3 Buffer Set	100 tests	(none)

### Product Notices

1. This reagent has been pre-diluted for use at the recommended Volume per Test. We typically use  $1 \times 10^6$  cells in a 100-μl experimental sample (a test).
2. Please refer to [www.bdbiosciences.com/pharmingen/protocols](http://www.bdbiosciences.com/pharmingen/protocols) for technical protocols.
3. An isotype control should be used at the same concentration as the antibody of interest.
4. For fluorochrome spectra and suitable instrument settings, please refer to our Fluorochrome Web Page at [www.bdbiosciences.com/colors](http://www.bdbiosciences.com/colors).
5. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
6. Source of all serum proteins is from USDA inspected abattoirs located in the United States.
7. Licensed for Research Purposes Only. Commercial use requires license from Boyce Thompson Institute for Plant Research.

### References

Fouser LA, Wright JF, Dunussi-Joannopoulos K, Collins M. Th17 cytokines and their emerging roles in inflammation and autoimmunity. *Immunol Rev.* 2008; 226:87-102. (Biology)

Shen F, Gaffen SL. Structure-function relationships in the IL-17 receptor: implications for signal transduction and therapy. *Cytokine.* 2008; 41(2):92-104. (Biology)

Starnes T, Robertson MJ, Sledge G, et al.. Cutting edge: IL-17F, a novel cytokine selectively expressed in activated T cells and monocytes, regulates angiogenesis and endothelial cell cytokine production. *J Immunol.* 2001; 167(8):4137-4140. (Biology)

Wang YH, Liu YJ. The IL-17 cytokine family and their role in allergic inflammation. *Curr Opin Immunol.* 2008; 20(6):697-702. (Biology)

Wright JF, Bennett F, Li B, et al. The human IL-17F/IL-17A heterodimeric cytokine signals through the IL-17RA/IL-17RC receptor complex. *J Immunol.* 2008; 181(4):2799-2805. (Biology)

Wright JF, Guo Y, Quazi A, et al. Identification of an interleukin 17F/17A heterodimer in activated human CD4+ T cells. *J Biol Chem.* 2007; 282(18):13447-13455. (Biology)