# Technical Data Sheet

# V450 Mouse IgA, κ Isotype Control

#### **Product Information**

**Material Number:** 562142

Alternate Name: IgA, κ; Immunoglobulin A, kappa

Size  $0.1 \, \text{mg}$ **Concentration:** 0.2 mg/ml M18-254 Clone:

Isotype: Mouse (BALB/c) IgA, κ Reactivity: QC Testing: Rat

Mouse and Human tested in development

Storage Buffer: Aqueous buffered solution containing protein stabilizer, glycerol and ≤0.09%

sodium azide.

## Description

The M18-254 antibody has unknown specificity and can be used as an immunoglobulin isotype control.

The antibody is conjugated to BD Horizon<sup>TM</sup> V450, which has been developed for use in multicolor flow cytometry experiments and is available exclusively from BD Biosciences. It is excited by the Violet laser Ex max of 406 nm and has an Em Max at 450 nm. Conjugates with BD Horizon<sup>TM</sup> V450 can be used in place of Pacific Blue<sup>TM</sup> conjugates.

## Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with BD Horizon™ V450 under optimum conditions, and unreacted BD Horizon™ V450 was removed.

#### **Application Notes**

# Application

Аррисации				
Flow cytometry	Routinely Tested			
Isotype control	Routinely Tested			

## **Suggested Companion Products**

Catalog Number	Name	Size	Clone
554656	Stain Buffer (FBS)	500 ml	(none)

### **Product Notices**

- Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
- Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
- For fluorochrome spectra and suitable instrument settings, please refer to our Fluorochrome Web Page at www.bdbiosciences.com/colors.
- BD Horizon<sup>TM</sup> V450 has a maximum absorption of 406 nm and maximum emission of 450 nm. Before staining with this reagent, please confirm that your flow cytometer is capable of exciting the fluorochrome and discriminating the resulting fluorescence.
- Pacific Blue™ is a trademark of Molecular Probes, Inc., Eugene, OR.
- An isotype control should be used at the same concentration as the antibody of interest.

# **BD Biosciences**

bdbiosciences.com

Asia Pacific Europe Japan 877.232.8995 888.268.5430 32.53.720.550 0120.8555.90 65.6861.0633 0800.771.7157

For country-specific contact information, visit bdbiosciences.com/how\_to\_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2011 BD

