

## Technical Data Sheet


## PE Hamster Anti-Rat CD29

## Product Information

<b>Material Number:</b>	<b>562154</b>
<b>Alternate Name:</b>	Itgb1; Integrin $\beta$ 1 chain; Integrin beta-1; VLA-4 subunit beta
<b>Size:</b>	50 $\mu$ g
<b>Concentration:</b>	0.2 mg/ml
<b>Clone:</b>	Ha2/5
<b>Immunogen:</b>	Rat glomerular epithelial cells
<b>Isotype:</b>	Armenian Hamster IgM, $\kappa$
<b>Reactivity:</b>	QC Testing: Rat
<b>Storage Buffer:</b>	Aqueous buffered solution containing protein stabilizer and $\leq 0.09\%$ sodium azide.

## Description

The Ha2/5 monoclonal antibody specifically binds to the 130 kDa integrin  $\beta$ 1 chain (CD29). CD29 is expressed on the cell surface as a heterodimer with one of the distinct integrin  $\alpha$  chains. With  $\alpha$ 1 through  $\alpha$ 6 (CD49a through CD49f), it forms the VLA-1 through VLA-6 complexes, respectively, and with  $\alpha$ V (CD51), it forms  $\alpha$ V $\beta$ 1 integrin. As a result, CD29 has a broad tissue distribution, including lymphocytes, endothelia, smooth muscle, and epithelia. The Ha2/5 hamster anti-rat CD29 monoclonal antibody cross-reacts with mouse thymocytes, splenocytes, and peripheral lymph node leukocytes. The Ha2/5 antibody blocks *in vitro* adhesion of CD29-expressing cells to collagen.



**Flow cytometric analysis of CD29 expression on rat splenocytes.** Lewis rat splenocytes were stained with either PE Armenian Hamster IgM,  $\lambda$ 1 Isotype Control (Cat. No. 562114, dashed line histogram) or a PE Hamster Anti-Rat CD29 antibody (Cat. No. 562154, solid line histogram). Flow cytometric fluorescence histograms were derived from gated events based on forward and side light-scatter characteristics of viable splenocytes. Flow cytometry was performed using a BD™ LSR II Flow Cytometer System.

## Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with R-PE under optimum conditions, and unconjugated antibody and free PE were removed.

## Application Notes

## Application

Flow cytometry	Routinely Tested
----------------	------------------

## Recommended Assay Procedure:

## Suggested Companion Products

Catalog Number	Name	Size	Clone
562114	PE Hamster IgM, $\lambda$ 1 Isotype Control	0.1 mg	G235-1
554656	Stain Buffer (FBS)	500 ml	(none)

## BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.268.5430	32.53.720.550	0120.8555.90	65.6861.0633	0800.771.7157

For country-specific contact information, visit [bdbiosciences.com/how\\_to\\_order/](http://bdbiosciences.com/how_to_order/)

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2011 BD



## Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to [www.bdbiosciences.com/pharmingen/protocols](http://www.bdbiosciences.com/pharmingen/protocols) for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
4. For fluorochrome spectra and suitable instrument settings, please refer to our Fluorochrome Web Page at [www.bdbiosciences.com/colors](http://www.bdbiosciences.com/colors).
5. Although hamster immunoglobulin isotypes have not been well defined, BD Biosciences Pharmingen has grouped Armenian and Syrian hamster IgG monoclonal antibodies according to their reactivity with a panel of mouse anti-hamster IgG mAbs. A table of the hamster IgG groups, Reactivity of Mouse Anti-Hamster Ig mAbs, may be viewed at [http://www.bdbiosciences.com/pharmingen/hamster\\_chart\\_11x17.pdf](http://www.bdbiosciences.com/pharmingen/hamster_chart_11x17.pdf).

## References

Mendrick DL, Kelly DM. Temporal expression of VLA-2 and modulation of its ligand specificity by rat glomerular epithelial cells in vitro. *Lab Invest.* 1993; 69(6):690-702. (Immunogen)

Springer TA. Adhesion receptors of the immune system. *Nature.* 1990; 346(6283):425-434. (Biology)