Technical Data Sheet

PE-CF594 Mouse Anti-Human IgM

Product Information


Material Number:							
Alternate Name:							
Size:							
Vol. per Test:							
Clone:							
Isotype:							
Reactivity:							
Storage Buffer:							

562539 IGHM; MU; Ig mu chain C region; VH 50 tests 5 µl G20-127 Mouse IgG1, ĸ QC Testing: Human Aqueous buffered solution containing BSA and ≤0.09% sodium azide.

Description

The G20-127 monoclonal antibody specifically binds to the heavy chain of human IgM. The majority of mature B lymphocytes express IgM. The G20-127 antibody does not react with other immunoglobulin heavy chain isotypes.

This antibody is conjugated to BD HorizonTM PE-CF594, which has been developed exclusively by BD Biosciences as a better alternative to PE-Texas Red®. PE-CF594 excites and emits at similar wavelengths to PE-Texas Red® yet exhibits improved brightness and spectral characteristics. Due to PE having maximal absorption peaks at 496 nm and 564 nm, PE-CF594 can be excited by the blue (488-nm), green (532-nm) and yellow-green (561-nm) lasers and can be detected with the same filter set as PE-Texas Red® (eg 610/20-nm filter).



Flow cytometric analysis of IgM expression on human peripheral blood lymphocytes. Human peripheral blood mononuclear cells were cultured in complete medium overnight in order to minimize subsequent nonspecific immunofluorescent staining. The cells were harvested and stained with BD Horizon™ PE-CF594 Mouse Anti-Human IgM (Cat. No. 562539) and APC Mouse Anti-Human CD19 (Cat. No. 555415/561742) antibodies. A two-color flow cytometric dot plot showing the correlated expression of cell surface IgM versus CD19 was derived from gated events with the forward and side light-scatter characteristics of viable lymphocytes. Flow cytometry was performed using a BD™ LSR II Flow Cytometer System.

Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. The antibody was conjugated with BD Horizon™ PE-CF594 under optimum conditions, and unconjugated antibody and free PE-CF594

Application Notes

were removed.

Flow cytometry	Routinely	Routinely Tested		
Suggested Compa	nion Products			
talog Number Name		Size	Clone	
54656	Stain Buffer (FBS)	500 ml	(none)	
55415	APC Mouse Anti-Human CD19	100 tests	HIB19	
61742	APC Mouse Anti-Human CD19	25 tests	HIB19	
62292	PE-CF594 Mouse IgG1, κ Isotype Control	0.1 mg	X40	

Product Notices

This reagent has been pre-diluted for use at the recommended Volume per Test. We typically use 1×10^{6} cells in a 100-µl experimental

sample (a test).

BD Biosciences

	bdbiosciences.com							
	United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean		
	877.232.8995	800.979.9408	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995		
For country contact information, visit bdbiosciences.com/contact								
Conditions: The information disclosed berein is not to be constructed as a recommendation to use the above produ								

Conditions: The information disclosed herein is not to be constructed as a recommendation to use the above product in violatio of any patents. BD Biosciences will not be help responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is stictly prohibited. For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale. Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD

- Source of all serum proteins is from USDA inspected abattoirs located in the United States. 2.
- 3. An isotype control should be used at the same concentration as the antibody of interest.
- 4. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
- Please observe the following precautions: Absorption of visible light can significantly alter the energy transfer occurring in any tandem 5. fluorochrome conjugate; therefore, we recommend that special precautions be taken (such as wrapping vials, tubes, or racks in aluminum foil) to prevent exposure of conjugated reagents, including cells stained with those reagents, to room illumination.
- Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before 6. discarding to avoid accumulation of potentially explosive deposits in plumbing.
- 7. For fluorochrome spectra and suitable instrument settings, please refer to our Multicolor Flow Cytometry web page at www.bdbiosciences.com/colors.
- Texas Red is a registered trademark of Molecular Probes, Inc., Eugene, OR. 8.
- 9. CF™ is a trademark of Biotium, Inc.
- When excited by the yellow-green (561-nm) laser, the fluorescence may be brighter than when excited by the blue (488-nm) laser. 10.
- This product is provided under an Agreement between BIOTIUM and BD Biosciences. The manufacture, use, sale, offer for sale, or import 11 of this product is subject to one or more patents or pending applications owned or licensed by Biotium, Inc. This product, and only in the amount purchased by buyer, may be used solely for buyer's own internal research, in a manner consistent with the accompanying product literature. No other right to use, sell or otherwise transfer (a) this product, or (b) its components is hereby granted expressly, by implication or by estoppel. This product is for research use only. Diagnostic uses require a separate license from Biotium, Inc. For information on purchasing a license to this product including for purposes other than research, contact Biotium, Inc., 3159 Corporate Place, Hayward, CA 94545, Tel: (510) 265-1027. Fax: (510) 265-1352. Email: btinfo@biotium.com.
- 12. Because of the broad absorption spectrum of the tandem fluorochrome, extra care must be taken when using multi-laser cytometers, which may directly excite both PE and CF™594.

References

Chtanova T, Tangye SG, Newton R, Frank N, Hodge MR, Rolph MS, Mackay CR. T follicular helper cells express a distinctive transcriptional profile, reflecting their role as non-Th1/Th2 effector cells that provide help for B cells. J Immunol. 2004; 173(1):68-78. (Clone-specific: Flow cytometry)

Gathings WE, Lawton AR, Cooper MD. Immunofluorescent studies of the development of pre-B cells, B lymphocytes and immunoglobulin isotype diversity in humans. Eur J Immunol. 1977; 7(11):804-810. (Biology)

Kuritani T, Cooper MD. Human B cell differentiation. II. Pokeweed mitogen-responsive B cells belong to a surface immunoglobulin D-negative subpopulation. J Exp Med. 1982: 155(5):1561-1566. (Biology)


Zola H, Macardle PJ, Flego L, Webster J. The expression of sub-population markers on B cells: a re-evaluation using high-sensitivity fluorescence flow cytometry. Dis Markers, 1991; 9(2);103-118, (Biology)

BD Biosciences

bdbiosciences.com United States Canada Europe 877.232.8995

Japan 800.979.9408 32.53.720.550 0120.8555.90 For country contact information, visit bdbiosciences.com/contact

Latin America/Caribbean 65.6861.0633 55.11.5185.9995



Asia Pacific

written authorization of Becton, Dickinson and Company is stictly prohibited. For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale. Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD

