

Technical Data Sheet


Alexa Fluor® 647 Mouse Anti-Human CD147

Product Information

Material Number:	562551
Alternate Name:	BSG; Basigin; Basi; Neurothelin; 5F7; EMMPRIN; M6; OK; TCSF
Size:	50 tests
Vol. per Test:	5 µl
Clone:	HIM6
Isotype:	Mouse IgG1, κ
Reactivity:	QC Testing: Human
Workshop:	VI NL109
Storage Buffer:	Aqueous buffered solution containing BSA and ≤0.09% sodium azide.

Description

The HIM6 monoclonal antibody specifically binds to CD147 which is encoded by BSG. CD147 is a type I transmembrane glycoprotein (30-50 kDa) of the immunoglobulin super-gene family. Neurothelin, a blood-brain barrier-specific molecule, was clustered as CD147 in the Sixth Human Leukocyte Differentiation Antigen (HLDA) workshop. It bears homology with mouse gp42 or basigin, human "M6" or "EMMPRIN", rat OX-47 or CD-9, and avian HT7 or 5A11. CD147 is also known as Tumor cell-derived collagenase stimulatory factor (TCSF). CD147 is a molecule that is broadly expressed on cells of hematopoietic and non-hematopoietic origin. Its expression on specific cell types may be regulated by cytokines. CD147 plays a role in embryonal blood-brain barrier development and a role in integrin-mediated adhesion in brain endothelia.



Flow cytometric analysis of CD147 expression on human peripheral blood lymphocytes. Whole blood was stained with Alexa Fluor® 647 Mouse Anti-Human CD147 antibody (Cat. No. 562551; solid line histogram) or with an Alexa Fluor® 647 Mouse IgG1, κ Isotype Control (Cat. No. 557714; dashed line histogram). Erythrocytes were lysed with BD Pharm Lyse™ Lysing Buffer (Cat. No. 555899). The fluorescence histograms were derived from gated events with the forward and side light-scatter characteristics of viable lymphocytes. Flow cytometry was performed using a BD™ LSR II Flow Cytometer System.

Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated to Alexa Fluor® 647 under optimum conditions, and unreacted Alexa Fluor® 647 was removed.

Application Notes

Application

Flow cytometry	Routinely Tested
----------------	------------------

Suggested Companion Products

Catalog Number	Name	Size	Clone
557714	Alexa Fluor® 647 Mouse IgG1 κ Isotype Control	100 tests	MOPC-21
555899	Lysing Buffer	100 ml	(none)
554656	Stain Buffer (FBS)	500 ml	(none)

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	800.979.9408	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD



Product Notices

1. This reagent has been pre-diluted for use at the recommended Volume per Test. We typically use 1×10^6 cells in a 100- μ l experimental sample (a test).
2. An isotype control should be used at the same concentration as the antibody of interest.
3. Source of all serum proteins is from USDA inspected abattoirs located in the United States.
4. Please refer to www.bdbiosciences.com/pharming/en/protocols for technical protocols.
5. The Alexa Fluor®, Pacific Blue™, and Cascade Blue® dye antibody conjugates in this product are sold under license from Molecular Probes, Inc. for research use only, excluding use in combination with microarrays, or as analyte specific reagents. The Alexa Fluor® dyes (except for Alexa Fluor® 430), Pacific Blue™ dye, and Cascade Blue® dye are covered by pending and issued patents.
6. Alexa Fluor® is a registered trademark of Molecular Probes, Inc., Eugene, OR.
7. Alexa Fluor® 647 fluorochrome emission is collected at the same instrument settings as for allophycocyanin (APC).
8. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
9. For fluorochrome spectra and suitable instrument settings, please refer to our Multicolor Flow Cytometry web page at www.bdbiosciences.com/colors.

References

Majdic O, Pickl WF, Kohl P, Stockinger H, Knapp W. EC16.3 CD147 Workshop: Reactivity and epitope mapping of CD147 monoclonal antibodies. In: Kishimoto T, Kikutani H, von dem Borne AEGK, ed. *Leukocyte Typing VI: White Cell Differentiation Antigens*. New York: Garland Publishing Inc; 1998:765-766. (Clone-specific: Flow cytometry)

Riethdorf S, Reimers N, Assmann V, Kornfeld JW, Terracciano L, Sauter G, Pantel K. High incidence of EMMPRIN expression in human tumors. *Int J Cancer*. 2006; 119(8):1800-1810. (Clone-specific: Immunohistochemistry, Immunoprecipitation, Western blot)

Rizzo A, Aragona E, Dino O, et al. EC16.1 CD147 Workshop: Expression of CD147 (neurothelin) in liver and lung cancer. In: Kishimoto T, Kikutani H, von dem Borne AEGK, ed. *Leukocyte Typing VI: White Cell Differentiation Antigens*. New York: Garland Publishing Inc; 1998:763-764. (Clone-specific: Immunohistochemistry)

Stockinger H, Ebel T, Hansmann C, et al. EC16 CD147 (neurothelin/basigin) Workshop Panel Report. In: Kishimoto T, Kikutani H, von dem Borne AEGK, ed. *Leukocyte Typing VI: White Cell Differentiation Antigens*. New York: Garland Publishing Inc; 1998:760-763. (Clone-specific: Flow cytometry, Immunoprecipitation)

Sudou A, Ozawa M, Muramatsu T. Lewis X structure increases cell substratum adhesion in L cells. *J Biochem (Tokyo)*. 1995; 117(2):271-275. (Biology)

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	800.979.9408	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD

