

Technical Data Sheet

PE-CF594 Mouse Anti-Human Bcl-2 (pS70)


Product Information

Material Number:	562679
Alternate Name:	BCL2; Apoptosis regulator Bcl-2; B-cell CLL/lymphoma 2; PPP1R50
Size:	50 tests
Vol. per Test:	5 µl
Clone:	N46-467
Immunogen:	Phosphorylated Peptide
Isotype:	Mouse IgG1
Reactivity:	QC Testing: Human Tested in Development: Mouse
Storage Buffer:	Aqueous buffered solution containing BSA and ≤0.09% sodium azide.

Description

The N46-467 monoclonal antibody specifically binds to Bcl-2 (pS70), ie, the Bcl-2 protein phosphorylated at the Ser70 site. Bcl-2 is a ~26 kDa intracellular, integral membrane protein found primarily in the nuclear envelope, endoplasmic reticulum and outer mitochondrial membrane. Bcl-2 is encoded by the *BCL2* (B-cell CLL/lymphoma 2) gene and is also known as Apoptosis regulator Bcl-2. Members of the Bcl-2 family play a major role in regulating the response of cells to apoptotic signals. Bcl-2 is one of the anti-apoptotic members of the Bcl-2 family. Bcl-2 knockout mice showed pronounced lymphoid apoptosis and other apoptosis related lesions later in life. Bcl-2 is a proto-oncogene because it blocks apoptosis and provides a selective survival advantage in many cell types and thus contributes to tumorigenesis. It has been implicated in several types of cancers, such as breast, prostate, and melanoma. Bcl-2 contains multiple phosphorylation sites including Thr56, Ser70, Thr74 and Ser87. Phosphorylation of Bcl-2 Ser70 has been shown to be a mitotic marker. Phosphorylation at this site regulates Bcl-2's anti-apoptotic activity and has recently been implicated in promoting autophagy. Several studies have shown that Bcl-2 phosphorylation is caused by c-Jun N-terminal kinase (JNK).

This antibody is conjugated to BD Horizon™ PE-CF594, which has been developed exclusively by BD Biosciences as a better alternative to PE-Texas Red®. PE-CF594 excites and emits at similar wavelengths to PE-Texas Red® yet exhibits improved brightness and spectral characteristics. Due to PE having maximal absorption peaks at 496 nm and 564 nm, PE-CF594 can be excited by the blue (488-nm), green (532-nm) and yellow-green (561-nm) lasers and can be detected with the same filter set as PE-Texas Red® (eg 610/20-nm filter).



Flow cytometric (Panels 1a and 1b) and Western blot (Panel 1c) analyses of Bcl-2 (pS70) expressed by Jurkat cells. Jurkat cells were either not treated (Panel 1a, dashed line histogram; Panel 1c, C) or treated (Panel 1a, solid line histogram; Panel 1b; Panel 1c, T) with 100 nM Taxol (Paclitaxel; Sigma, Cat. No. T7191; 24 hr, 37°C). For Panel 1a, cells were fixed in BD Phosflow™ Cytotfix Buffer (Cat. No. 554655; 10 min, 37°C) and permeabilized in BD Phosflow™ Perm Buffer III (Cat. No. 558050; 30 min, on ice) prior to staining with BD Phosflow™ PE-CF594 Mouse Anti-Bcl-2 (pS70) (Cat. No. 562679). For optimal co-staining of total cellular DNA (Panel 1b), cells were fixed and permeabilized in 70% ethanol (30 min, on ice) prior to staining with DAPI (Sigma, Cat. No. D8417) and PE-CF594 Mouse Anti-Bcl-2 (pS70). Histograms (Panel 1a) and a two-color dot plot showing DNA (DAPI) versus Bcl-2 (pS70) levels (Panel 1b) were generated from gated events with the light scatter characteristics of intact cells using a BD™ LSR II Flow Cytometer System. For Western blot (Panel 1c), cell lysates (15 µg total cell protein/lane) were blotted using Purified Mouse Anti-Bcl-2 (pS70) (Cat. No. 562529; 0.125, 0.063, 0.032 µg/ml for Lanes 1, 2, and 3, respectively). Bcl-2 (pS70) was identified as a ~28 kDa band.

Flow cytometric (Panels 2a and 2b) and Western blot (Panel 2c) analyses of Bcl-2 (pS70) expressed by human peripheral blood mononuclear cells. PHA-stimulated (Sigma Cat. No. L1668; 20 µg/ml for 3 days) PBMC were either not treated (Panel 2a, dashed line histogram; Panel 2c, C) or treated (Panel 2a, solid line histogram; Panel 2b; Panel 2c, T) with Taxol (100 nM, 24 hr, 37°C). Flow cytometric analyses of Bcl-2 (pS70) expression without (Panel 2a) or with (Panel 2b) co-staining for DNA content were performed using a BD LSRFortessa™ Cell Analyzer System. Lysates from 1X10⁶ PBMC were blotted using Purified Mouse Anti-Bcl-2 (pS70) antibody (2.0 µg/ml) as described above.

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	800.979.9408	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD



Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with BD Horizon™ PE-CF594 under optimum conditions, and unconjugated antibody and free PE-CF594 were removed.

Application Notes

Application

Intracellular staining (flow cytometry)

Routinely Tested

Purified or conjugated mAb were characterized by flow cytometry (Flow), Western blot (WB), and immunohistochemistry (IHC) using these model systems:

Method	Species	Cells	Treatment	Fixation	Perm buffer	Result
Flow	Human	PHA-stimulated PBMC	Nocodazole	Cytofix	Perm III	Induced in a subpopulation of cells
	Human	PHA-stimulated PBMC	Taxol	Cytofix	Perm III	Induced in a subpopulation of cells
	Human	Jurkat (serum-starved)	Taxol	Cytofix	Perm III	Induced in most cells. Blocked by pS70 phospho peptide but not by non-phospho peptide.
	Human	PBMC	PMA	Cytofix	Perm III	Weakly induced
WB	Human	PHA-stimulated PBMC	Nocodazole			28-kDa band increased
	Human	PHA-stimulated PBMC	Taxol			28-kDa band increased
	Human	Jurkat (serum-starved)	Taxol			28-kDa band increased. Blocked by pS70 phospho peptide but not by non-phospho peptide.
	Human	PBMC	PMA			28-kDa band increased
IHC	Human	Tonsil	Formalin fixed human paraffin tonsil sections with citrate buffer pretreatment			Cytoplasmic and nuclear staining observed

Suggested Companion Products

Catalog Number	Name	Size	Clone
554655	Fixation Buffer	100 ml	(none)
558050	Perm Buffer III	125 ml	(none)
554656	Stain Buffer (FBS)	500 ml	(none)
562529	Purified Mouse anti-Human Bcl-2 (pS70)	0.1 mg	N46-467

Product Notices

1. This reagent has been pre-diluted for use at the recommended Volume per Test. We typically use 1×10^6 cells in a 100- μ l experimental sample (a test).
2. Source of all serum proteins is from USDA inspected abattoirs located in the United States.
3. Please refer to www.bdbiosciences.com/pharming/en/protocols for technical protocols.
4. Please observe the following precautions: Absorption of visible light can significantly alter the energy transfer occurring in any tandem fluorochrome conjugate; therefore, we recommend that special precautions be taken (such as wrapping vials, tubes, or racks in aluminum foil) to prevent exposure of conjugated reagents, including cells stained with those reagents, to room illumination.
5. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
6. For fluorochrome spectra and suitable instrument settings, please refer to our Multicolor Flow Cytometry web page at www.bdbiosciences.com/colors.
7. Texas Red is a registered trademark of Molecular Probes, Inc., Eugene, OR.
8. CF™ is a trademark of Biotium, Inc.
9. When excited by the yellow-green (561-nm) laser, the fluorescence may be brighter than when excited by the blue (488-nm) laser.
10. This product is provided under an Agreement between BIOTIUM and BD Biosciences. The manufacture, use, sale, offer for sale, or import of this product is subject to one or more patents or pending applications owned or licensed by Biotium, Inc. This product, and only in the amount purchased by buyer, may be used solely for buyer's own internal research, in a manner consistent with the accompanying product literature. No other right to use, sell or otherwise transfer (a) this product, or (b) its components is hereby granted expressly, by implication or by estoppel. This product is for research use only. Diagnostic uses require a separate license from Biotium, Inc. For information on purchasing a license to this product including for purposes other than research, contact Biotium, Inc., 3159 Corporate Place, Hayward, CA 94545, Tel: (510) 265-1027. Fax: (510) 265-1352. Email: btinfo@biotium.com.
11. Because of the broad absorption spectrum of the tandem fluorochrome, extra care must be taken when using multi-laser cytometers, which may directly excite both PE and CF™594.
12. Species testing during development may have been performed with a different format of the same clone. Selected applications have been tested for cross-reactivity.
13. All other brands are trademarks of their respective owners.

References

Deng X, Xiao L, Lang W, Gao F, Ruvolo P, May WS, Jr.. Novel role for JNK as a stress-activated Bcl2 kinase. *J Biol Chem*. 2001; 276(26):23681-23688. (Biology)
Geng F, Tang L, Li Y, Yang L, Choi KS, Kazim AL, Zhang Y.. Allyl isothiocyanate arrests cancer cells in mitosis, and mitotic arrest in turn leads to apoptosis via Bcl-2 protein phosphorylation. *J Biol Chem*. 2011; 286(37):32259-32267. (Biology)
Ling YH, Tornos C, Perez-Soler R. Phosphorylation of Bcl-2 is a marker of M phase events and not a determinant of apoptosis. *J Biol Chem*. 1998; 273(30):18984-18991. (Biology)

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	800.979.9408	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD



Maundrell K, Antonsson B, Magnenat E, Camps M, Muda M, Chabert C, Gillieron C, Boschert U, Vial-Knecht E, Martinou JC, Arkinstall S.. Bcl-2 undergoes phosphorylation by c-Jun N-terminal kinase/stress-activated protein kinases in the presence of the constitutively active GTP-binding protein Rac1. *J Biol Chem.* 1997; 272(40):25238-42. (Biology)

Pattingre S, Bauvy C, Carpentier S, Levade T, Levine B, Codogno P.. Role of JNK1-dependent Bcl-2 phosphorylation in ceramide-induced macroautophagy. *J Biol Chem.* 2009; 284(5):2719-2728. (Biology)

Sarkar S, Korolchuk VI, Renna M, Imarisio S, Fleming A, Williams A, Garcia-Arencibia M, Rose C, Luo S, Underwood BR, Kroemer G, O'Kane CJ, Rubinsztein. Complex inhibitory effects of nitric oxide on autophagy. *Mol Cell.* 2011; 43(1):19-32. (Biology)

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	800.979.9408	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be constructed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD

