Technical Data Sheet

BV421 Mouse Anti-SSEA-1

Product Information

Material Number: Alternate Name: Size Vol. per Test: **Clone:** Isotype: **Reactivity:**


562705 3-FAL, X-hapten, LeX antigen, CD15 50 tests 5 µl MC480 Mouse IgM, ĸ QC Testing: Mouse Reported: Human Aqueous buffered solution containing BSA and ≤0.09% sodium azide.

Storage Buffer:

Description

The MC480 monoclonal antibody reacts with Stage-Specific Embryonic Antigen-1 (SSEA-1), which is a terminal carbohydrate epitope (3-fucosyl-N-acetyllactosamine or 3-FAL) on glycoproteins and lactose series glycolipids. SSEA-1 is related to Lewis blood group antigens and is found in a variety of embryonic and adult tissues and cancers. As its name implies, the expression of SSEA-1 is stage-specific and can be used to characterize embryonic cells and monitor their differentiation. However, its expression pattern differs in the human and mouse. In the human, SSEA-1 is not found on embryonic stem (ES) cells, embryonic inner cell mass (ICM), or teratocarcinoma (embryonal carcinoma or EC) cells. As human EC and ES cells undergo differentiation, SSEA-1 expression is upregulated. In the adult, the same epitope is expressed as CD15 on granulocytes and monocytes, but not lymphocytes or dendritic cells. In the mouse, SSEA-1 is found on EC, ES, and primordial germ cells, 8-cell to blastocyst embryos, ICM, and on subpopulations of cells in the adult central nervous system, including stem cells. In contrast to the human, SSEA-1 expression is reduced as mouse EC and ES cells undergo differentiation.

The antibody was conjugated to BD Horizon[™] BV421 which is part of the BD Horizon[™] Brilliant Violet[™] family of dyes. With an Ex Max of 407-nm and Em Max at 421-nm, BD HorizonTM BV421 can be excited by the violet laser and detected in the standard Pacific BlueTM filter set (eg, 450/50-nm filter). BD HorizonTM BV421 conjugates are very bright, often exhibiting a 10 fold improvement in brightness compared to Pacific BlueTM conjugates.



Flow cytometric analysis of SSEA-1 expression on mouse embryonic stem (ES) cells. ES-E14TG2a mouse ES cells (ATCC, CRL-1821™) at passage 36 were harvested with Accutase™ (Cat. No. 561527) and stained with BD Horizon™ BV421 Mouse Anti-SSEA-1 (solid line) or a BD Horizon™ BV421 Mouse IgM, κ isotype control (Cat. No.562704, dashed line). Flow cvtometry was performed on a BD™ LSRII flow cytometry system.

Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with BD HorizonTM BV421 under optimum conditions, and unconjugated antibody and free BD HorizonTM BV421 were removed.

Application Notes

Application Flow cytometry Routinely Tested **BD Biosciences** bdbiosciences.com United States Canada Europe Japan Asia Pacific Latin America/Caribbean 877.232.8995 800.979.9408 32.53.720.550 0120.8555.90 65.6861.0633 55.11.5185.9995 For country contact information, visit bdbiosciences.com/contact Conditions: The information disclosed herein is not to be constructed as a recommendation to use the above product in violation

of any patents. BD Biosciences will not be help responsible for patent infringement or other violations that may occur with the use of our products. Buckness will not be help responsible for patent infringement or other violations that may occur with the product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is stictly prohibited. For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale. Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD

Suggested Companion Products

Catalog Number	Name	Size	Clone
561527	Accutase [™] Cell Detachment Solution	100 ml	(none)
562704	BV421 Mouse IgM, κ Isotype Control	50 µg	G155-228
554656	Stain Buffer (FBS)	500 ml	(none)

Product Notices

- This reagent has been pre-diluted for use at the recommended Volume per Test. We typically use 1×10^{6} cells in a 100-µl experimental 1. sample (a test).
- 2. An isotype control should be used at the same concentration as the antibody of interest.
- Brilliant Violet[™] 421 is a trademark of Sirigen. 3
- Accutase is a registered trademark of Innovative Cell Technologies, Inc. 4.
- 5. Pacific Blue[™] is a trademark of Molecular Probes, Inc., Eugene, OR.
- Source of all serum proteins is from USDA inspected abattoirs located in the United States. 6.
- 7. Caution: Sodium azide vields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
- 8. For fluorochrome spectra and suitable instrument settings, please refer to our Multicolor Flow Cytometry web page at www.bdbiosciences.com/colors.
- 9. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.

References

Capela A, Temple S. LeX/ssea-1 is expressed by adult mouse CNS stem cells, identifying them as nonependymal. Neuron. 2002; 35:865-875. (Biology) Childs RA, Pennington J, Uemura K, et al. High-molecular-weight glycoproteins are the major carriers of the carbohydrate differentiation antigens I, i and SSEA-1 of mouse teratocarcinoma cells. Biochem J. 1983; 215:491-503. (Clone-specific)

Draper JS, Pigott C, Thomson JA, Andrews PW. Surface antigens of human embryonic stem cells: changes upon differentiation in culture. J Anat. 2002; 200:249-258. (Clone-specific)

Henderson JK, Draper JS, Baillie HS, et al. Preimplantation human embryos and embryonic stem cells show comparable expression of stage-specific embryonic antigens. Stem Cells. 2002; 20:329-337. (Clone-specific)

Kannagi R, Nudelman E, Levery SB, Hakomori S. A series of human erythrocyte glycosphingolipids reacting to the monoclonal antibody directed to a developmentally regulated antigen, SSEA-1. J Biol Chem. 1982; 257(24):14865-14874. (Clone-specific)

Solter D, Knowles BB. Monoclonal antibody defining a stage-specific mouse embryonic antigen (SSEA-1). Proc Natl Acad Sci U S A. 1978; 75(11):5565-5569. (Immunogen)


Thomson JA, Itskovitz-Eldor J, Shapiro SS, et al. Embryonic stem cell lines derived from human blastocysts. Science. 1998; 282:1145-1147. (Clone-specific)

BD Biosciences

bdbiosciences.com United States Canada 877.232.8995

Europe Japan 800.979.9408 32.53.720.550 0120.8555.90 For country contact information, visit bdbiosciences.com/contact

Latin America/Caribbean 65.6861.0633 55.11.5185.9995



Asia Pacific

written authorization of Becton, Dickinson and Company is stictly prohibited. For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale. Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD