

Technical Data Sheet


BV421 Mouse Anti-Human TGF-β1**Product Information**

Material Number:	562962
Alternate Name:	TGFB1; TGFbeta1; TGF-beta-1; Transforming growth factor, beta 1; CED; LAP
Size:	50 tests
Vol. per Test:	5 µl
Clone:	TW4-9E7
Immunogen:	Human TGF-β1 Transfected Cell Line
Isotype:	Mouse (BALB/c) IgG1, κ
Reactivity:	QC Testing: Human
Storage Buffer:	Aqueous buffered solution containing BSA and ≤0.09% sodium azide.

Description

The TW4-9E7 monoclonal antibody specifically binds to Human Transforming Growth Factor beta-1 (TGF-β1). TGF-β1 is a potent multifunctional cytokine that positively or negatively regulates numerous processes including development, hematopoiesis, tissue remodeling, wound repair, innate and adaptive immunity as well as cancer and autoimmune diseases. TGF-β1 is formed by the enzymatic cleavage of the TGF-β1 propeptide that is encoded by the *TGFB1* gene and comprised of the Latency Associated Peptide (LAP) and TGF-β1. Prior to secretion, the dimeric LAP-TGF-β1 propeptide is cleaved resulting in a biologically inactive form of dimeric TGF-β1 that is noncovalently associated with dimeric LAP (latent TGF-β1). This complex may be expressed on the surface of TGF-β1-producing cells or be further processed by proteolytic removal of LAP to release the biologically active mature form of the soluble TGF-β1 homodimer. Many different cell types synthesize TGF-β1 and express specific receptors for it. The TW4-9E7 antibody recognizes both the intracellular latent bound form of TGF-β1 along with the membrane bound form of TGF-β1.

The antibody was conjugated to BD Horizon™ BV421 which is part of the BD Horizon™ Brilliant Violet™ family of dyes. With an Ex Max of 407-nm and Em Max at 421-nm, BD Horizon™ BV421 can be excited by the violet laser and detected in the standard Pacific Blue™ filter set (eg, 450/50-nm filter). BD Horizon™ BV421 conjugates are very bright, often exhibiting a 10 fold improvement in brightness compared to Pacific Blue™ conjugates.



Flow cytometric analysis of human TGF-β1 expressed by TGF-β1-transfected P3UI cells. Untransfected mouse P3UI myeloma cells (dashed line histogram) and human TGF-β1-transfected P3UI cells (solid line histogram) were fixed and permeabilized for 30 minutes with BD Cytofix/Cytoperm™ Fixation and Permeabilization Solution (Cat. No. 554722) and washed with BD Perm/Wash™ Perm/Wash Buffer (Cat. No. 554723). The cells were then stained with BD Horizon™ BV421 Mouse Anti-Human TGF-β1 antibody (Cat. No. 562962). The flow cytometric fluorescence histograms were derived from gated events with the forward- and side light-scattering characteristics of intact cells. Flow cytometry was performed using a BD™ LSR II Flow Cytometer System.

Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with BD Horizon™ BV421 under optimum conditions, and unconjugated antibody and free BD Horizon™ BV421 were removed.

Application Notes**Application**

Intracellular staining (flow cytometry)	Routinely Tested
---	------------------

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	800.979.9408	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.
For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.
Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD



Suggested Companion Products

Catalog Number	Name	Size	Clone
554656	Stain Buffer (FBS)	500 ml	(none)
562438	BV421 Mouse IgG1, k Isotype Control	50 µg	X40
554722	Fixation and Permeabilization Solution	125 ml	(none)
554723	Perm/Wash Buffer	100 ml	(none)

Product Notices

1. This reagent has been pre-diluted for use at the recommended Volume per Test. We typically use 1×10^6 cells in a 100-µl experimental sample (a test).
2. Source of all serum proteins is from USDA inspected abattoirs located in the United States.
3. An isotype control should be used at the same concentration as the antibody of interest.
4. Please refer to www.bdbiosciences.com/pharming/protocols for technical protocols.
5. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
6. For fluorochrome spectra and suitable instrument settings, please refer to our Multicolor Flow Cytometry web page at www.bdbiosciences.com/colors.
7. Pacific Blue™ is a trademark of Molecular Probes, Inc., Eugene, OR.
8. Brilliant Violet™ 421 is a trademark of Sirigen.

References

Derynck R, Akhurst R.J. Differentiation plasticity regulated by TGF-beta family proteins in development and disease. *Nat Cell Biol.* 2007; 9(9):1000-1004. (Biology)

Derynck R, Jarrett JA, Chen EY, et al. Human transforming growth factor-beta complementary DNA sequence and expression in normal and transformed cells. *Nature.* 1985; 316(6030):701-705. (Biology)

Dünker N, Kriegelstein K. Targeted mutations of transforming growth factor-beta genes reveal important roles in mouse development and adult homeostasis. *Eur J Biochem.* 2000; 267(24):6982-6988. (Biology)

Miyazono K, Hellman U, Wernstedt C, Heldin CH. Latent high molecular weight complex of transforming growth factor beta 1. Purification from human platelets and structural characterization. *J Biol Chem.* 1988; 263(13):6407-6415. (Biology)

Oida T, Weiner HL. Overexpression of TGF-β1 gene induces cell surface localized glucose-regulated protein 78-associated latency-associated peptide/TGF-β. *J Immunol.* 2010; 185(6):3529-3535. (Clone-specific: Flow cytometry)

Rubtsov YP, Rudensky AY. TGFbeta signalling in control of T-cell-mediated self-reactivity. *Nat Rev Immunol.* 2007; 7(6):443-453. (Biology)

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	800.979.9408	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD

