# **Technical Data Sheet**

# BV650 Mouse Anti-Human CD335 (NKp46)

### **Product Information**

**Material Number:** 563230

Alternate Name: NCR1; NK-p46; hNKp46; LY94; Natural cytotoxicity triggering receptor 1

Size Vol. per Test: 5 μl 9E2/Nkp46 Clone:

Human NKp46 Recombinant Protein Immunogen:


Isotype: Mouse (BALB/c) IgG1, κ Reactivity: QC Testing: Human

Storage Buffer: Aqueous buffered solution containing BSA and ≤0.09% sodium azide.

# Description

The 9E2/Nkp46 monoclonal antibody specifically binds to CD335. CD335 is also known as the Natural killer cell p46-related protein (NKp46) and the Natural cytotoxicity triggering receptor 1 (NCR1). CD335 is a 46 kDa type I membrane glycoprotein that is expressed on resting and activated NK cells. Its extracellular region contains two C2-type, Ig-like domains. The transmembrane domain contains a positively charged amino acid (Arg) which could be involved in stabilizing the association with CD3 $\zeta$ . Its intracellular region does not contain immunoreceptor tyrosine-based activating motifs (ITAM), but it is linked to intracytoplasmic transduction machinery by its association with CD3\(\zeta\) and Fc\(\text{RI}\) adaptor proteins. CD335 along with NKp30 and NKp44 are referred to as natural cytotoxicity receptors (NCR). These receptors play very important roles in cells that kill virus-infected target cells, tumor cells and MHC-class I-unprotected cells.

The antibody was conjugated to BD Horizon™ BV650 which is part of the BD Horizon™ Brilliant Violet™ family of dyes. This dye is a tandem fluorochrome of BD Horizon™ BV421 with an Ex Max of 405-nm and an acceptor dye with an Em Max at 650-nm. BD Horizon™ BV650 can be excited by the violet laser and detected in a filter used to detect APC-like dyes (eg, 660/20-nm filter). Due to the excitation and emission characteristics of the acceptor dye, there will be spillover into the APC and Alexa Fluor® 700 detectors. However, the spillover can be corrected through compensation as with any other dye combination.



Flow cytometric analysis of CD335 (NKp46) expression on human peripheral blood lymphocytes. Whole blood was stained with BD Horizon™ BV650 Mouse Anti-Human CD335 (NKp46) antibody (Cat. No. 563230; solid line histogram) or with a BD Horizon™ BV650 Mouse IgG1, κ Isotype Control (Cat. No. 563231; dashed line histogram) Erythrocytes were lysed with BD PharmLyse™ Lysing Buffer (Cat. No. 555899). The fluorescence histograms were derived from gated events with the forward and side light-scatter characteristics of viable lymphocytes. Flow cytometry was performed using a BD™ LSR II Flow Cytometer System.

# Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with BD Horizon<sup>TM</sup> BV650 under optimum conditions, and unconjugated antibody and free BD Horizon<sup>TM</sup> BV650 were removed.

## **Application Notes**

Application

Flow cytometry Routinely Tested

# **BD Biosciences**

bdbiosciences.com

United States Canada Europe Asia Pacific Latin America/Caribbean 877.232.8995 800.979.9408 32.53.720.550 0120.8555.90 65.6861.0633 55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be constructed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be help responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is stictly prohibited.
For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.
Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD



# **Suggested Companion Products**

| Catalog Number | Name                                | Size   | Clone  |  |
|----------------|-------------------------------------|--------|--------|--|
| 554656         | Stain Buffer (FBS)                  | 500 ml | (none) |  |
| 563231         | BV650 Mouse IgG1, k Isotype Control | 50 μg  | X40    |  |
| 555899         | Lysing Buffer                       | 100 ml | (none) |  |

#### **Product Notices**

- This reagent has been pre-diluted for use at the recommended Volume per Test. We typically use 1 × 10<sup>6</sup> cells in a 100-µl experimental sample (a test).
- 2. An isotype control should be used at the same concentration as the antibody of interest.
- 3. Brilliant Violet<sup>TM</sup> 650 is a trademark of Sirigen.
- 4. Alexa Fluor® is a registered trademark of Molecular Probes, Inc., Eugene, OR.
- 5. Source of all serum proteins is from USDA inspected abattoirs located in the United States.
- Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
- For fluorochrome spectra and suitable instrument settings, please refer to our Multicolor Flow Cytometry web page at www.bdbiosciences.com/colors.
- 8. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.

#### References

Mandelboim O, Porgador A. NKp46. Int J Biochem Cell Biol. 2001; 33(12):1147-1150. (Biology)

Nakajima H, Cella M, Bouchon A, et al. Patients with X-linked lymphoproliferative disease have a defect in 2B4 receptor-mediated NK cell cytotoxicity. *Eur J Immunol.* 2000; 30(11):3309-3318. (Immunogen: Flow cytometry)

Sivori S, Pende D, Bottino C, et al. NKp46 is the major triggering receptor involved in the natural cytotoxicity of fresh or cultured human NK cells. Correlation between surface density of NKp46 and natural cytotoxicity against autologous, allogeneic or xenogeneic target cells. *Eur J Immunol.* 1999; 29(5):1656-1666. (Biology)

Sivori S, Vitale M, Morelli L, et al. p46, a novel natural killer cell-specific surface molecule that mediates cell activation. *J Exp Med.* 1997; 186(7):1129-1136. (Biology)

# **BD Biosciences**

bdbiosciences.com

 United States
 Canada
 Europe
 Japan
 Asia Pacific
 Latin America/Caribbean

 877.232.8995
 800.979.9408
 32.53.720.550
 0120.8555.90
 65.6861.0633
 55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be constructed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be help responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is stictly prohibited. For Research Use Only, Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD



563230 Rev. 1 Page 2 of 2