# **Technical Data Sheet**

# BV650 Rat Anti-Human CD197 (CCR7)

# **Product Information**

Material Number:	563407
Alternate Name:	CCR7, BLR-2, EBI-1, CMKBR7
Size:	50 tests
Vol. per Test:	5 µl
Clone:	3D12
Immunogen:	Human CCR7 protein
Isotype:	Rat IgG2a, ĸ
Reactivity:	QC Testing: Human
Storage Buffer:	Aqueous buffered solution containing BSA and ≤0.09% sodium azide.

### Description

The monoclonal antibody 3D12 reacts with the human CC chemokine receptor, CCR7. CCR7 (previously known as BLR-2, EBI-1 and CMKBR7), a seven-transmembrane, G-protein-coupled receptor, is the specific receptor for CC chemokines, MIP-3 $\beta$ /Exodus 3/ELC/ CCL19 and 6Ckine/Exodus 2/SLC/TCA4/CCL21. It has been shown that CCR7 mRNA is expressed mainly in lymphoid tissues including spleen, lymph nodes and tonsil. CCR7 mRNA was also detected in peripheral T and B lymphocytes, in bone marrow and cord blood CD34-positive cells and mature dendritic cells. The human CCR7 gene, unlike other CC chemokine receptor genes, has been mapped to chromosome 17q12. The immunogen used to generate 3D12 hybridoma was the N-terminus as well as parts of the second extracellular loop of human CCR7 protein. The monoclonal antibody 3D12 recognizes an epitope mapping to the N-terminus of human CCR7.

The antibody was conjugated to BD Horizon<sup>™</sup> BV650 which is part of the BD Horizon<sup>™</sup> Brilliant Violet<sup>™</sup> family of dyes. This dye is a tandem fluorochrome of BD Horizon<sup>™</sup> BV421 with an Ex Max of 405-nm and an acceptor dye with an Em Max at 650-nm. BD Horizon<sup>™</sup> BV650 can be excited by the violet laser and detected in a filter used to detect APC-like dyes (eg, 660/20-nm filter). Due to the excitation and emission characteristics of the acceptor dye, there will be spillover into the APC and Alexa Fluor® 700 detectors. However, the spillover can be corrected through compensation as with any other dye combination.



Multicolor flow cytometric analysis of CD197 (CCR7) expression on CD4+ and CD8+ human peripheral blood lymphocytes. Human whole blood was stained with FITC Mouse Anti-Human CD45RA (Cat. No. 555488/561882) and BD Horizon™ BV650 Rat Anti-Human CD197 (CCR7) (Cat. No. 563407) and either PerCP-CY™5.5 Mouse Anti-Human CD4 (Cat. No. 560650; Left Panel) or Alexa Fluor® 647 Mouse Anti-Human CD8 (Cat. No. 557708; Right Panel). Erythrocytes were lysed with BD Pharm Lyse™ Lysing Buffer (Cat. No. 555899). The two-color flow cytometric dot plots show the correlated expression patterns of CD45RA versus CD197 (CCR7) for CD4+ or CD8+ gated events with the forward and side light-scatter characteristics of viable lymphocytes. Flow cytometric analysis was performed using a BD™ LSR II Flow Cytometer System.

# **Preparation and Storage**

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with BD Horizon<sup>™</sup> BV650 under optimum conditions, and unconjugated antibody and free BD Horizon<sup>™</sup>

BV650 were removed.

BD Biosciences							
bdbiosciences.com							
United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean		
877.232.8995	800.979.9408	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995		
For country contact information, visit bdbiosciences.com/contact							
Conditions: The ir of any patents. Bl use of our produce product or as a co written authoriza	nformation disclose D Biosciences will n ts. Purchase does n Imponent of anoth tion of Becton, Dic	d herein is not to b ot be help responsi ot include or carry er product. Any use kinson and Compar	e constructed as a n ble for patent infrin any right to resell o of this product oth ny is stictly prohibite	ecommendation to u gement or other vic r transfer this produ ter than the permitte ed.	use the above product in violation plations that may occur with the ct either as a stand-alone ed use without the express		
For Research Use	Only. Not for use in	diagnostic or there	apeutic procedures.	Not for resale.			
Unless otherwise	noted, BD, BD Logo	and all other trade	emarks are property	of Becton, Dickinso	n and Company. © 2011 BD		



#### **Application Notes**

Application

Flow cytometry Routinely Tested

# Suggested Companion Products

Catalog Number	Name	Size	Clone	
554656	Stain Buffer (FBS)	500 ml	(none)	
563236	BV650 Rat IgG2a, κ Isotype Control	50 µg	R35-95	
555899	Lysing Buffer	100 ml	(none)	
555488	FITC Mouse Anti-Human CD45RA	100 tests	HI100	
561882	FITC Mouse Anti-Human CD45RA	25 tests	HI100	
560650	PerCP-Cy <sup>™</sup> 5.5 Mouse Anti-Human CD4	50 tests	RPA-T4	
557708	Alexa Fluor® 647 Mouse Anti-Human CD8	100 tests	RPA-T8	

#### **Product Notices**

- 1. This reagent has been pre-diluted for use at the recommended Volume per Test. We typically use  $1 \times 10^{6}$  cells in a 100-µl experimental sample (a test).
- 2. Source of all serum proteins is from USDA inspected abattoirs located in the United States.
- An isotype control should be used at the same concentration as the antibody of interest. 3.
- 4. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
- Alexa Fluor® is a registered trademark of Molecular Probes, Inc., Eugene, OR. 5.
- Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before 6. discarding to avoid accumulation of potentially explosive deposits in plumbing.
- 7. For fluorochrome spectra and suitable instrument settings, please refer to our Multicolor Flow Cytometry web page at www.bdbiosciences.com/colors.
- Brilliant Violet<sup>™</sup> 650 is a trademark of Sirigen. 8.

#### References

Birkenbach M, Josefsen K, Yalamanchili R, Lenoir G, Kieff E. Epstein-Barr virus-induced genes: first lymphocyte-specific G protein-coupled peptide receptors. Nature. 1993; 67(4):2209-2220. (Biology)

Burgstahler R, Kempkes B, Steube K, Lipp M. Expression of the chemokine receptor BLR2/EBI1 is specifically transactivated by Epstein-Barr virus nuclear antigen 2. Biochem Biophys Res Commun. 1995; 215(2):737-743. (Biology)

Kim CH, Pelus LM, White JR, Broxmeyer HE. Macrophage-inflammatory protein-3 beta/EBI1-ligand chemokine/CK beta-11, a CC chemokine, is a

chemoattractant with a specificity for macrophage progenitors among myeloid progenitor cells. J Immunol. 1998; 161(5):2580-2585. (Biology)

Lipp M, Burgstahler R, Muller G, et al. Functional organization of secondary lymphoid organs by the chemokine system. Curr Top Microbiol Immunol. 2000; 251:173-179. (Immunogen: Functional assay, Inhibition)

Sallusto F, Lenig D, Forster R, Lipp M, Lanzavecchia A. Two subsets of memory T lymphocytes with distinct homing potentials and effector functions. Nature. 1999; 401(6754):708-712. (Clone-specific: Flow cytometry)

Schweickart VL, Raport CJ, Godiska R, et al. Cloning of human and mouse EBI1, a lymphoid-specific G-protein-coupled receptor encoded on human chromosome 17q12-q21.2. Genomics. 1994; 23(3):643-650. (Biology)

Yanagihara S, Komura E, Nagafune J, Watarai H, Yanaguchi Y. EBI1/CCR7 is a new member of dendritic cell chemokine receptor that is up-regulated upon maturation. J Immunol. 1998; 161(6):3096-3102. (Biology)

Yoshida R, Imai T, Hieshima K, et al. Molecular cloning of a novel human CC chemokine EBI1-ligand chemokine that is a specific functional ligand for EBI1, CCR7. J Biol Chem. 1997; 272(21):13803-13809. (Biology)

Yoshida R, Nagira M, Imai T, et al. EBI1-ligand chemokine (ELC) attracts a broad spectrum of lymphocytes: activated T cells strongly up-regulate CCR7 and efficiently migrate toward ELC. Int Immunol. 1998; 10(7):901-910. (Biology)

Yoshida R, Nagira M, Kitaura M, Imagawa N, Imai T, Yoshie O. Secondary lymphoid-tissue chemokine is a functional ligand for the CC chemokine receptor CCR7. J Biol Chem. 1998; 273(12):7118-7122. (Biology)

#### **BD Biosciences**

bubiosciences.	com	
United States	Canada	Europe
877.232.8995	800.979.9408	32.53.720
-		

.550 0120.8555.90 65.6861.0633 For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be constructed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be help responsible for patent infringement or other violations that may occur with the use of our products. Buckness will not be help responsible for patent infringement or other violations that may occur with the product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is stictly prohibited. For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale. Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD

Asia Pacific

Latin America/Caribbean

55.11.5185.9995

Japan

