

## Technical Data Sheet


**BV510 Mouse Anti-Human CD16****Product Information**

<b>Material Number:</b>	<b>563830</b>
<b>Alternate Name:</b>	FcRIII; Fc-gamma RIII; FCG3; FCGR3; FCGRIII; FcγRIII; IGFR3
<b>Size:</b>	100 tests
<b>Vol. per Test:</b>	5 µl
<b>Clone:</b>	3G8
<b>Immunogen:</b>	Human polymorphonuclear leukocytes
<b>Isotype:</b>	Mouse (BALB/c x DBA/2) IgG1, κ
<b>Reactivity:</b>	QC Testing: Human Tested in Development: Rhesus, Cynomolgus, Baboon
<b>Workshop:</b>	IV N409
<b>Storage Buffer:</b>	Aqueous buffered solution containing BSA and ≤0.09% sodium azide.

**Description**

The 3G8 monoclonal antibody specifically binds to the 50-65 kDa transmembrane form of the IgG Fc Receptor (FcγRIII), a human NK cell-associated antigen. CD16 is expressed on NK cells as well as macrophages and granulocytes. Reports indicate that CD16 plays a role in signal transduction and NK cell activation. The 3G8 antibody blocks the binding of soluble immune complexes to granulocytes. The 3G8 antibody is reported (Vossebeld *et al.*, 1997) to increase intracellular calcium levels in human neutrophils by interacting with both FcγRIIIa and FcγRIIIb molecules. This antibody has also been reported to induce homotypic neutrophil aggregation.

The antibody was conjugated to BD Horizon™ BV510 which is part of the BD Horizon™ Brilliant Violet™ family of dyes. With an Ex Max of 405-nm and Em Max at 510-nm, BD Horizon™ BV510 can be excited by the violet laser and detected in the BD Horizon™ V500 (525/50-nm) filter set. BD Horizon™ BV510 conjugates are useful for the detection of dim markers off the violet laser.



**Two-color flow cytometric analysis of human CD16 expression on human peripheral blood cells.** Human peripheral blood cells were stained with APC Mouse Anti-Human CD56 antibody (Cat. No. 555518) and either BD Horizon™ BV510 Mouse IgG1, κ Isotype Control (Cat. No. 562946; Left Panel) or BD Horizon™ BV510 Mouse Anti-Human CD16 antibody (Cat. No. 563829/563830; Right Panel). The erythrocytes were lysed with BD FACS™ Lysing Solution (Cat. No. 349202). The two-color flow cytometric dot plots show the correlated expression patterns of CD16 versus CD56 (or Ig Isotype control staining) for gated events with the forward and side light-scatter characteristics of intact peripheral blood lymphocytes. Flow cytometric analysis was performed using a BD™ LSR II Flow Cytometer System.

**Preparation and Storage**

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with BD Horizon™ BV510 under optimum conditions, and unconjugated antibody and free BD Horizon™ BV510 were removed.

**BD Biosciences**

[bdbiosciences.com](http://bdbiosciences.com)

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	800.979.9408	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country contact information, visit [bdbiosciences.com/contact](http://bdbiosciences.com/contact)

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD



## Application Notes

### Application

Flow cytometry

Routinely Tested

### Suggested Companion Products

Catalog Number	Name	Size	Clone
554656	Stain Buffer (FBS)	500 ml	(none)
562946	BV510 Mouse IgG1, k Isotype Control	50 µg	X40
563829	BV510 Mouse Anti-Human CD16	25 tests	3G8
555518	APC Mouse Anti-Human CD56	100 tests	B159
349202	BD FACST <sup>™</sup> Lysing Solution	100 ml	(none)
555899	Lysing Buffer	100 ml	(none)

### Product Notices

1. This reagent has been pre-diluted for use at the recommended Volume per Test. We typically use  $1 \times 10^6$  cells in a 100-µl experimental sample (a test).
2. An isotype control should be used at the same concentration as the antibody of interest.
3. Species testing during development may have been performed with a different format of the same clone. Selected applications have been tested for cross-reactivity.
4. Brilliant Violet<sup>™</sup> 510 is a trademark of Sirigen.
5. Source of all serum proteins is from USDA inspected abattoirs located in the United States.
6. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
7. For fluorochrome spectra and suitable instrument settings, please refer to our Multicolor Flow Cytometry web page at [www.bdbiosciences.com/colors](http://www.bdbiosciences.com/colors).
8. Please refer to [www.bdbiosciences.com/pharming/protocols](http://www.bdbiosciences.com/pharming/protocols) for technical protocols.

### References

Fleit HB, Wright SD, Unkeless JC. Human neutrophil Fc gamma receptor distribution and structure. *Proc Natl Acad Sci U S A*. 1982; 79(10):3275-3279. (Immunogen: Blocking, Immunoprecipitation, Inhibition, Radioimmunoassay)

Knapp W, Dörken B, Gilks WR, et al, ed. *Leucocyte Typing IV*. New York, NY: Oxford University Press; 1989:1-1182. (Clone-specific)

Perussia B, Trinchieri G, Jackson A, et al. The Fc receptor for IgG on human natural killer cells: phenotypic, functional, and comparative studies with monoclonal antibodies. *J Immunol*. 1984; 133(1):180-189. (Clone-specific: Flow cytometry, Functional assay, Inhibition)

Schmidt RE. Non-lineage/natural killer section report: new and previously defined clusters. In: Knapp W, Dörken B, Gilks WR, et al, ed. *Leucocyte Typing IV: White Cell Differentiation Antigens*. New York, NY: Oxford University Press; 1989:517-542. (Clone-specific)

Stroncek DF, Skubitz KM, Plachta LB, et al. Alloimmune neonatal neutropenia due to an antibody to the neutrophil Fc-gamma receptor III with maternal deficiency of CD16 antigen. *Blood*. 1991; 77(7):1572-1580. (Clone-specific: Immunofluorescence, Immunoprecipitation)

Vossebeld PJ, Homburg CH, Roos D, Verhoeven AJ. The anti-Fc gamma RIII mAb 3G8 induces neutrophil activation via a cooperative action of Fc gamma RIIIb and Fc gamma RIIa. *Int J Biochem Cell Biol*. 1997; 29(3):465-473. (Clone-specific: Activation, Functional assay)

Wirthmueller U, Kurosaki T, Murakami MS, Ravetch JV. Signal transduction by Fc gamma RIII (CD16) is mediated through the gamma chain. *J Exp Med*. 1992; 175(5):1381-1390. (Clone-specific: Activation, Functional assay)

Zola H, Swart B, Nicholson I, Voss E. *Leukocyte and Stromal Cell Molecules. The CD Markers*. Hoboken, New Jersey: John Wiley & Sons, Inc.; 2007:1-581. (Biology)

### BD Biosciences

[bdbiosciences.com](http://bdbiosciences.com)

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	800.979.9408	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country contact information, visit [bdbiosciences.com/contact](http://bdbiosciences.com/contact)

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD

