

Technical Data Sheet


Purified Mouse Anti-Csk

Product Information


Material Number:	610079
Size:	50 µg
Concentration:	250 µg/ml
Clone:	52/Csk
Immunogen:	Rat Csk aa. 1-156
Isotype:	Mouse IgG1
Reactivity:	QC Testing: Mouse Tested in Development: Human, Dog, Rat, Chicken
Target MW:	50 kDa
Storage Buffer:	Aqueous buffered solution containing BSA, glycerol, and ≤0.09% sodium azide.

Description

The *csk* gene encodes a protein kinase that phosphorylates several Src family kinases including p60[c-src], p59[fyn], p53/56[lyn], p56[lck], and p62[yes]. This phosphorylation occurs specifically at the carboxy-terminal tyrosine, Y527. Csk (c-src kinase) was identified based on its ability to phosphorylate exogenous Src in mammalian *src*-deficient cells. Establishment of a mouse *csk*- cell line revealed that phosphorylation of Y527 in Src was reduced 20-50%. This correlated with an increase of Src kinase activity, thus indicating Csk is a negative regulator of *src*. *Csk*-deficient cells also show an elevation in the kinase activity of p59[fyn] and p53/56[lyn]. These data suggest that Csk may be a universal negative regulator of Src family kinases in vivo.



Western blot analysis of Csk on mouse macrophage lysate. Lane 1: 1:500, lane 2: 1:1000, lane 3: 1:2000 dilution of anti-Csk.



Immunohistochemical staining of Rabbit Spleen.

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

Store undiluted at -20° C.

Application Notes

Application

Western blot	Routinely Tested
Immunoprecipitation	Tested During Development
Immunofluorescence	Tested During Development
Immunohistochemistry	Tested During Development

BD Biosciences

bdbiosciences.com

United States 877.232.8995 Canada 888.259.0187 Europe 32.53.720.550 Japan 0120.8555.90 Asia Pacific 65.6861.0633 Latin America/Caribbean 55.11.5185.9995

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2008 BD



Suggested Companion Products

Catalog Number	Name	Size	Clone
611479	Mouse Macrophage Cell Lysate	500 µg	(none)
554002	HRP Goat Anti-Mouse Ig	1.0 ml	(none)
554001	FITC Goat Anti-Mouse Ig	0.5 mg	Polyclonal

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to www.bdbiosciences.com/pharming/en/protocols for technical protocols.
3. Source of all serum proteins is from USDA inspected abattoirs located in the United States.
4. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

References

Bénistant C, Bourgaux JF, Chapuis H, Mottet N, Roche S, Bali JP. The COOH-terminal Src kinase Csk is a tumor antigen in human carcinoma. *Cancer Res.* 2001; 61(4):1415-1420.(Clone-specific: Immunofluorescence, Western blot)

Bergman M, Mustelin T, Oetken C. The human p50csk tyrosine kinase phosphorylates p56lck at Tyr-505 and down regulates its catalytic activity. *EMBO J.* 1992; 11(8):2919-2924.(Biology)

Okada M, Nada S, Yamanashi Y, Yamamoto T, Nakagawa H. CSK: a protein-tyrosine kinase involved in regulation of src family kinases. *J Biol Chem.* 1991; 266(36):24249-24252.(Biology)

Takayama Y, Tanaka S, Nagai K, Okada M. Adenovirus-mediated overexpression of C-terminal Src kinase (Csk) in type I astrocytes interferes with cell spreading and attachment to fibronectin. Correlation with tyrosine phosphorylations of paxillin and FAK. *J Biol Chem.* 1999; 274(4):2291-2297.(Clone-specific: Immunoprecipitation)

Wan Y, Bence K, Hata A, Kurosaki T, Veillette A, Huang XY. Genetic evidence for a tyrosine kinase cascade preceding the mitogen-activated protein kinase cascade in vertebrate G protein signaling. *J Biol Chem.* 1999; 272(27):17209-17215.(Clone-specific: In vitro kinase assay, Western blot)