

Technical Data Sheet


Purified Mouse Anti-CD95

Product Information


Material Number: 610198
Alternate Name: Fas, APO-1
Size: 150 µg
Concentration: 250 µg/ml
Clone: 13/Fas
Immunogen: Human Fas aa. 1-163
Isotype: Mouse IgG2a, κ
Reactivity: QC Testing: Human
 Tested in Development: Mouse, Rat, Dog, Chicken
Target MW: 45 kDa
Storage Buffer: Aqueous buffered solution containing BSA, glycerol, and ≤0.09% sodium azide.

Description

CD95 is a member of a family of cell surface receptors that includes tumor necrosis factor receptor (TNF-R), nerve growth factor receptor (NGF-R), CD40, CD27, CD30, and 4-1BB. Both murine and human Fas genes have been cloned and reportedly share 60% similarity in their amino acid sequences. CD95 (Fas) is a cell surface apoptosis-signaling molecule that is widely expressed in sites such as thymus, liver, heart, and ovary. Abnormalities in the Fas gene correlate with autoimmune features in mice and with unusually high levels of lymphocyte apoptosis in HIV-infected humans. Genetic studies have localized the Fas gene near the lpr (lymphoproliferation disease) locus on mouse chromosome 19 and further characterization reportedly has revealed that lpr is a mutation affecting the function of the Fas gene.



Western blot analysis of CD95 on a Daudi cell lysate (Human B lymphoblast; ATCC CCL-213). Lane 1:5,000, lane 2: 1:10,000, lane 3: 1:20,000 in dilution of the Purified Mouse anti-CD95 antibody.



Immunohistochemical staining of a formalin-fixed rabbit liver section.

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. Store undiluted at -20°C.

Application Notes

Application

Western blot	Routinely Tested
Immunohistochemistry-formalin (antigen retrieval required)	Tested During Development
Immunofluorescence	Not Recommended
Immunoprecipitation	Not Recommended

BD Biosciences

bdbiosciences.com

United States 877.232.8995 Canada 800.979.9408 Europe 32.53.720.550 Japan 0120.8555.90 Asia Pacific 65.6861.0633 Latin America/Caribbean 55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be help responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD



Recommended Assay Procedure:

For Western Blot and immunohistochemistry resources, please reference <http://www.bdbiosciences.com/resources/cellbiology/index.jsp>

Suggested Companion Products

<u>Catalog Number</u>	<u>Name</u>	<u>Size</u>	<u>Clone</u>
554002	HRP Goat Anti-Mouse Ig	1.0 ml	(none)

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to www.bdbiosciences.com/pharming/en/protocols for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
4. Source of all serum proteins is from USDA inspected abattoirs located in the United States.

References

Arnold R, Seifert M, Asadullah K, Volk HD. Crosstalk between keratinocytes and T lymphocytes via Fas/Fas ligand interaction: modulation by cytokines.. *J Immunol.* 1999; 162(12):7140-7147. (Biology: Flow cytometry)

Itoh N, Yonehara S, Ishii A, et al. The polypeptide encoded by the cDNA for human cell surface antigen Fas can mediate apoptosis. *Cell.* 1991; 66(2):233-243. (Biology)

MacLaren A, Clark W, Gillespie DA. v-Jun sensitizes cells to apoptosis by a mechanism involving mitochondrial cytochrome C release.. *Oncogene.* 2000; 19(51):5906-5918. (Biology: Western blot)

Rosen K, Coll ML, Li A, Filmus J. Transforming growth factor-alpha prevents detachment-induced inhibition of c-Src kinase activity, Bcl-XL down-regulation, and apoptosis of intestinal epithelial cells.. 2001; 276(40):37273-37279. (Clone-specific: Western blot)

Zhuang S, Demirs JT, Kochevar IE. Protein kinase C inhibits singlet oxygen-induced apoptosis by decreasing caspase-8 activation.. *Oncogene.* 2001; 20(46):6764-6776. (Biology: Immunofluorescence, Immunoprecipitation, Western blot)

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	800.979.9408	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD

