

## Technical Data Sheet


**Purified Mouse Anti-Gephyrin****Product Information**

<b>Material Number:</b>	<b>610585</b>
<b>Size:</b>	150 µg
<b>Concentration:</b>	250 µg/ml
<b>Clone:</b>	45/Gephyrin
<b>Immunogen:</b>	Rat Gephyrin aa. 569-726
<b>Isotype:</b>	Mouse IgG1
<b>Reactivity:</b>	QC Testing: Rat Tested in Development: Chicken, Human, Mouse
<b>Target MW:</b>	93 kDa
<b>Storage Buffer:</b>	Aqueous buffered solution containing BSA, glycerol, and ≤0.09% sodium azide.


**Description**

The sub-membrane region of the postsynaptic neuron is an intricate network of cytoskeletal elements generally known as the postsynaptic density. It is thought that this elaborate cytoskeletal region is critical for receptor targeting, clustering, and efficient signal input. Gephyrin, a 93 kDa protein, was identified as a result of its ability to bind to polymerized tubulin (microtubules). Although expressed in all tissues, *gephyrin* mRNA is found most abundantly in brain. Alternative splicing produces at least five different mRNAs. Gephyrin colocalizes and interacts with the glycine receptor at the postsynaptic density. It is possible that Gephyrin acts as an anchor between the glycine receptor and microtubules.

This antibody is routinely tested by western blot analysis. Other applications were tested at BD Biosciences Pharmingen during antibody development only or reported in the literature.



**Western blot analysis of Gephyrin on rat brain lysate.**  
Lane 1: 1:250, lane 2: 1:500, lane 3: 1:1000 dilution of anti-Gephyrin.



**Rat cerebellum, formalin-fixed paraffin embedded tissue, with citrate pre-treatment, 20X.**

**Preparation and Storage**

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. Store undiluted at -20° C.

**BD Biosciences**

[www.bdbiosciences.com](http://www.bdbiosciences.com)

<b>United States</b>	<b>Canada</b>	<b>Europe</b>	<b>Japan</b>	<b>Asia Pacific</b>	<b>Latin America/Caribbean</b>
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country-specific contact information, visit [www.bdbiosciences.com/how\\_to\\_order/](http://www.bdbiosciences.com/how_to_order/)

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2007 BD



## Application Notes

### Application

Western blot	Routinely Tested
Immunoprecipitation	Tested During Development
Immunofluorescence	Tested During Development
Immunohistochemistry	Tested During Development

## Suggested Companion Products

Catalog Number	Name	Size	Clone
611464	Rat Cerebellum Lysate	500 µg	(none)
554001	FITC Goat Anti-Mouse Igs (Multiple Adsorption)	0.5 mg	Polyclonal
554002	HRP Goat Anti-Mouse Igs	1.0 ml	(none)

## Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to [www.bdbiosciences.com/pharming/protocols](http://www.bdbiosciences.com/pharming/protocols) for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
4. Source of all serum proteins is from USDA inspected abattoirs located in the United States.

## References

Feng G, Tintrup H, Kirsch J, et al. Dual requirement for gephyrin in glycine receptor clustering and molybdoenzyme activity. *Science*. 1998; 282(5392):1321-1324. (Clone-specific: Western blot)

Kneussel M, Haverkamp S, Fuhrmann JC, et al. The gamma-aminobutyric acid type A receptor (GABAAR)-associated protein GABARAP interacts with gephyrin but is not involved in receptor anchoring at the synapse. *Proc Natl Acad Sci U S A*. 2000; 97(15):8594-8599. (Clone-specific: Immunofluorescence, Western blot)

Meyer G, Kirsch J, Betz H, Langosch D. Identification of a gephyrin binding motif on the glycine receptor beta subunit. *Neuron*. 1995; 15(3):563-572. (Biology)

Prior P, Schmitt B, Grenningloh G, et al. Primary structure and alternative splice variants of gephyrin, a putative glycine receptor-tubulin linker protein. *Neuron*. 1992; 8(6):1161-1170. (Biology)

## BD Biosciences

[www.bdbiosciences.com](http://www.bdbiosciences.com)

United States 877.232.8995 Canada 888.259.0187 Europe 32.53.720.550 Japan 0120.8555.90 Asia Pacific 65.6861.0633 Latin America/Caribbean 55.11.5185.9995

For country-specific contact information, visit [www.bdbiosciences.com/how\\_to\\_order/](http://www.bdbiosciences.com/how_to_order/)

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2007 BD

