

Technical Data Sheet


Purified Mouse Anti-DGK θ

Product Information


Material Number:	610931
Size:	150 μ g
Concentration:	250 μ g/ml
Clone:	24/DGKtheta
Immunogen:	Human DGK θ aa. 677-883
Isotype:	Mouse IgG1
Reactivity:	QC Testing: Rat Tested in Development: Human, Mouse, Rabbit
Target MW:	110 kDa
Storage Buffer:	Aqueous buffered solution containing BSA, glycerol, and \leq 0.09% sodium azide.

Description

The protein kinase C pathway is a major signal transduction system that is activated upon stimulation of transmembrane receptors by hormones, neurotransmitters, and growth factors. Key mediators in this pathway are increased intracellular free Ca²⁺ levels and formation of diacylglycerol (DAG). DGK θ (diacylglycerol kinase θ) restricts PKC activation through the phosphorylation of DAG molecules that contain an unsaturated fatty acid at the sn-2 position to produce phosphatidic acid (PA). DGK θ contains several regions that are found in signaling molecules where they function in lipid-protein and protein-protein interactions. A C-terminal catalytic domain, three CRDs (cysteine rich domains), a PH domain, and an N-terminal proline/glycine rich domain are structural features of DGK θ . Six potential PKC phosphorylation sites lie between CRD3 and the PH domain. Cell-specific expression differentiate multiple isoforms of DGK. DGK θ is expressed primarily within the cerebellar cortex and hippocampus of the brain, but is also found in the small intestine and liver. The presence of the RA (Ras-associating) domain suggests that DGK θ may mediate activity of the Ras-like small GTP binding proteins.



Western blot analysis of DGK θ on rat brain lysate.
Lane 1: 1:250, lane 2: 1:500, lane 3: 1:1000 dilution of anti-DGK θ .



Immunofluorescent staining of SK-BR-3 cells.

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. Store undiluted at -20° C.

BD Biosciences

www.bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country-specific contact information, visit www.bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2007 BD



Application Notes

Application

Western blot	Routinely Tested
Immunofluorescence	Tested During Development
Immunohistochemistry	Tested During Development

Suggested Companion Products

Catalog Number	Name	Size	Clone
611463	Rat Cerebrum Lysate	500 µg	(none)
554002	HRP Goat Anti-Mouse Ig	1.0 ml	(none)
554001	FITC Goat Anti-Mouse Ig	0.5 mg	Polyclonal

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to www.bdbiosciences.com/pharming/en/protocols for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
4. Source of all serum proteins is from USDA inspected abattoirs located in the United States.

References

- Bregoli L, Baldassare JJ, Raben DM. Nuclear diacylglycerol kinase-theta is activated in response to alpha-thrombin. *J Biol Chem.* 2001; 276(26):23288-23295. (Clone-specific: Immunofluorescence, Western blot)
- Houssa B, Schaap D, van der Wal J, et al. Cloning of a novel human diacylglycerol kinase (DGKtheta) containing three cysteine-rich domains, a proline-rich region, and a pleckstrin homology domain with an overlapping Ras-associating domain. *J Biol Chem.* 1997; 272(16):10422-10428.(Biology)
- Pilz A, Schaap D, Hunt D, Fitzgibbon J. Chromosomal localization of three mouse diacylglycerol kinase (DAGK) genes: genes sharing sequence homology to the Drosophila retinal degeneration A (rdgA) gene. *Genomics.* 1995; 26(3):599-601.(Biology)