Technical Data Sheet

Purified Mouse Anti-α-Spectrin II

Product Information

Material Number: Size: Concentration: Clone: Immunogen: Isotype: Reactivity:

Target MW: Storage Buffer: 612560 50 μg 250 μg/ml 35/α-Spectrin II Human α-Spectrin II aa. 252-371 Mouse IgG1 QC Testing: Human Reactivity Confirmed in Development: Dog, Chicken Lack of Reactivity Confirmed in Development: Rat, Mouse 250 kDa Aqueous buffered solution containing BSA, glycerol, and ≤0.09% sodium azide.

Description

Spectrins are central components of the cytoskeleton that form a scaffold below the plasma membrane. Spectrins contain two subunits, α and β , which intertwine to form heterodimers that can self associate into elongated tetramers. α -spectrin I and β -spectrin I form heterodimers in red blood cells, while nonerythroid mammalian cells contain heterodimers of α -spectin I and II with β -spectrin I to V. The structure of spectrins includes a succession of triple-helical repeats along with various domains, such as SH3 domain, EF hands, PH domains, and binding domains for ankyrin, actin, band 4.1, and calmodulin. α -spectrin II is a widely expressed non-erythroid α -spectrin Hat contains an SH3 domain, a calmodulin binding site, and two cleavage sites for proteases, such as calpains and caspase-3. β -spectrin II is a widely expressed non-erythroid β -spectrin II and β -spectrin II, like many other spectrins, can form heterodimers that can self associate into tetramers, as well as interact with Band 4.1, F-actin, and other proteins near the plasma membrane. This scaffold of cytoskeletal and plasma membrane proteins is critical for the maintenance of cell structure.





E

Immunofluorescent staining of MDCK cells.

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. Store undiluted at -20°C.

Application Notes

Application

Western blot	Routinely Tested	
Immunofluorescence	Tested During Development	

Recommended Assay Procedure:

Western blot: Please refer to http://www.bdbiosciences.com/pharmingen/protocols/Western_Blotting.shtml.

BD Biosciences

bdbiosciences.	com				
United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995
For country-spe	ecific contact in	formation, visit	bdbiosciences.co	m/how_to_orde	r/
Conditions: The ir	nformation disclose	d herein is not to b	e construed as a rec	ommendation to use	e the above product in violation
of any patents. Bl	D Biosciences will n	ot be held responsi	ble for patent infrin	gement or other vio	lations that may occur with the
use of our produc	ts. Purchase does n	ot include or carry	any right to resell o	r transfer this produ	ct either as a stand-alone
product or as a co	omponent of anoth	er product. Any us	e of this product oth	ner than the permitt	ed use without the express
written authoriza	tion of Becton Dick	kinson and Compan	y is strictly prohibite	ed.	
For Research Use	Only. Not for use in	diagnostic or there	peutic procedures.	Not for resale.	
BD, BD Logo and	all other trademark	ks are the property	of Becton, Dickinsor	n and Company. ©20	108 BD



Suggested Companion Products

Catalog Number	Name	Size	Clone
611451	Jurkat Cell Lysate	500 μg	(none)
554002	HRP Goat Anti-Mouse Ig	1.0 ml	(none)
554001	FITC Goat Anti-Mouse Ig	0.5 mg	Polyclonal

Product Notices

- 1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
- 2. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
- 3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
- 4. Source of all serum proteins is from USDA inspected abattoirs located in the United States.

References

Hu RJ, Watanabe M, Bennett V. Characterization of human brain cDNA encoding the general isoform of beta-spectrin. J Biol Chem. 1992; 267(26):18715-18722. (Biology)

Moon RT, McMahon AP. Generation of diversity in nonerythroid spectrins. Multiple polypeptides are predicted by sequence analysis of cDNAs encompassing the coding region of human nonerythroid alpha-spectrin. *J Biol Chem.* 1990; 265(8):4427-4433.(Biology)

Nicolas G, Fournier CM, Galand C, et al. Tyrosine phosphorylation regulates alpha II spectrin cleavage by calpain. Mol Cell Biol. 2002; 22(10):3527-3536. (Biology)