

Technical Data Sheet

Purified Mouse Anti-mDia1

Product Information


| | |
|-------------------------|--|
| Material Number: | 610849 |
| Alternate Name: | p140mDia; Drf1 |
| Size: | 150 µg |
| Concentration: | 250 µg/ml |
| Clone: | 51/mDia1 |
| Immunogen: | Mouse p140mDia aa. 41-153 |
| Isotype: | Mouse IgG1 |
| Reactivity: | QC Testing: Mouse Tested in Development: Rat, Human, Dog |
| Target MW: | 140 kDa |
| Storage Buffer: | Aqueous buffered solution containing BSA, glycerol, and ≤0.09% sodium azide. |

Description

GTPases like Ras, Rho, cdc42Hs, and Rac modulate a multitude of cellular functions like cytoskeletal architecture, growth, motility, and gene expression. The activity of the GTP-binding proteins is regulated by factors that accelerate GTP-binding (GAPs) and proteins that enhance the rate of GTP hydrolysis. mDia1, also known as p140mDia and Drf1, is the mammalian homologue of *Drosophila's* diaphanous, a protein essential for cytokinesis. The 1255 amino acid mDia1 is widely expressed, with a Rho binding domain at the NH2-terminal region, a central polyproline region, and an FH2 domain. This protein also shares homology with the yeast Bn1p essential for budding and the mouse formin necessary for proper limb development. mDia1 binds to GTP-Rho and to the actin-binding protein profilin—all three proteins are co-localized at the lamellipodia in cultured cells. The overexpression of mDia1 promoted the formation of actin filaments, implicating this protein in cell motility events regulated by Rho. mDia1 was also found consistently mutated in familial deafness.



Western blot analysis of mDia1 on a RSV-3T3 cell lysate. Lane of 1: 1:500, lane 2: 1:1000, lane 3: 1:2000 dilution of the mouse anti-mDia1 antibody.



Immunofluorescence staining of FHS cells (Normal human fetal lung fibroblasts; ATCC HTB-157).

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

Store undiluted at -20°C.

Application Notes

Application

| | |
|----------------------|---------------------------|
| Western blot | Routinely Tested |
| Immunofluorescence | Tested During Development |
| Immunoprecipitation | Not Recommended |
| Immunohistochemistry | Not Recommended |

BD Biosciences

bdbiosciences.com

United States 877.232.8995 Canada 888.259.0187 Europe 32.53.720.550 Japan 0120.8555.90 Asia Pacific 65.6861.0633 Latin America/Caribbean 55.11.5185.9995

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2008 BD



Recommended Assay Procedure:

Western blot: Please refer to http://www.bdbiosciences.com/pharmingen/protocols/Western_Blotting.shtml

Suggested Companion Products

| <u>Catalog Number</u> | <u>Name</u> | <u>Size</u> | <u>Clone</u> |
|-----------------------|-------------------------|-------------|--------------|
| 554002 | HRP Goat Anti-Mouse Ig | 1.0 ml | (none) |
| 554001 | FITC Goat Anti-Mouse Ig | 0.5 mg | Polyclonal |

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
4. Source of all serum proteins is from USDA inspected abattoirs located in the United States.

References

Li F, Higgs HN. The mouse Formin mDia1 is a potent actin nucleation factor regulated by autoinhibition. *Curr Biol.* 2003; 13(15):1335-1340.(Biology)
Pennisi E. The architecture of hearing. *Science.* 1997; 278(5341):1223-1224.(Biology)
Watanabe N, Madaule P, Reid T. p140mDia, a mammalian homolog of Drosophila diaphanous, is a target protein for Rho small GTPase and is a ligand for profilin. *EMBO J.* 1997; 16(11):3044-3056.(Biology)