

## Technical Data Sheet

**Purified Mouse Anti-Human Btk****Product Information**

<b>Material Number:</b>	611117
<b>Size:</b>	150 µg
<b>Concentration:</b>	250 µg/ml
<b>Clone:</b>	53/BTK
<b>Immunogen:</b>	Human N-Terminal Btk aa. 2-172 Recombinant Protein
<b>Isotype:</b>	Mouse IgG2a
<b>Reactivity:</b>	QC Testing: Human
<b>Target MW:</b>	77 kDa
<b>Storage Buffer:</b>	Aqueous buffered solution containing BSA, glycerol, and ≤0.09% sodium azide.

**Description**

Bruton's tyrosine kinase (Btk) is a nonreceptor tyrosine kinase whose function is critical for proper B cell development and signaling. It is a member of the Tec family of kinases which includes Tec and Itk. This family is similar to the src family of tyrosine kinases. However, Tec family members lack the N-terminal myristylation site and the regulatory C-terminal tyrosine that are found in src proteins. In addition to an N-terminal pleckstrin homology (PH) domain, the Tec proteins contain Src homology domains 2 and 3 (SH2 and SH3) and a stretch of 60-80 amino acids between the PH and SH3 domains termed the Tec homology domain. The activity of Btk is regulated by Src-mediated phosphorylation of the kinase domain at tyrosine 551. This event induces Btk kinase activity and subsequent autophosphorylation at tyrosine 223 in the SH3 domain. Phosphorylated Btk then associates with the cell membrane via the interaction of the PH domain with phosphatidylinositol 3, 4, 5-triphosphate. The PH domain is essential for proper activation and function of Btk. A mutation in the PH domain results in Xid, murine X-linked immunodeficiency, and human X-linked agammaglobulinemia.



**Western blot analysis of Btk on EB1 lysate. Lane 1:**  
1:250, lane 2: 1:500, lane 3: 1:1000 dilution of anti-Btk.

**Preparation and Storage**

Store undiluted at -20° C.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

**Application Notes****Application**

Western blot	Routinely Tested
Immunofluorescence	Tested During Development

**BD Biosciences**

[www.bdbiosciences.com](http://www.bdbiosciences.com)

United States 877.232.8995    Canada 888.259.0187    Europe 32.53.720.550    Japan 0120.8555.90    Asia Pacific 65.6861.0633    Latin America/Caribbean 55.11.5185.9995

For country-specific contact information, visit [www.bdbiosciences.com/how\\_to\\_order/](http://www.bdbiosciences.com/how_to_order/)

*Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.*

*For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.*

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2007 BD



## Suggested Companion Products

Catalog Number	Name	Size	Clone
611546	EB1 Cell Lysate	500 µg	(none)
554001	FITC Goat Anti-Mouse Ig	0.5 mg	Polyclonal
554002	HRP Goat Anti-Mouse Ig	1.0 ml	(none)

## Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to [www.bdbiosciences.com/pharming/en/protocols](http://www.bdbiosciences.com/pharming/en/protocols) for technical protocols.
3. Source of all serum proteins is from USDA inspected abattoirs located in the United States.
4. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

## References

- Aoki Y, Isselbacher KJ, Pillai S. Bruton tyrosine kinase is tyrosine phosphorylated and activated in pre-B lymphocytes and receptor-ligated B cells. *Proc Natl Acad Sci U S A*. 1994; 91(22):10606-10609.(Biology)
- Shahan TA, Sorenson WG, Simpson J, Kefalides NA, Lewis DM. Tyrosine kinase activation in response to fungal spores is primarily dependent on endogenous reactive oxygen production in macrophages. *J Biol Chem*. 2000; 275(14):10175-10181.(Clone-specific: Immunoprecipitation, In vitro kinase assay, Western blot)
- Sideras P, Müller S, Shiels H. Genomic organization of mouse and human Bruton's agammaglobulinemia tyrosine kinase (Btk) loci. *J Immunol*. 1994; 153(12):5607-5617.(Biology)
- Vetrie D, Vorechovský I, Sideras P. The gene involved in X-linked agammaglobulinaemia is a member of the src family of protein-tyrosine kinases. *Nature*. 1993; 361(6409):226-233.(Biology)
- Yang W, Malek SN, Desiderio S. An SH3-binding site conserved in Bruton's tyrosine kinase and related tyrosine kinases mediates specific protein interactions in vitro and in vivo. *J Biol Chem*. 1995; 270(35):20832-20840.(Biology)