

## Technical Data Sheet


**Purified Mouse Anti-Mouse Nedd4****Product Information**

<b>Material Number:</b>	<b>611480</b>
<b>Size:</b>	50 µg
<b>Concentration:</b>	250 µg/ml
<b>Clone:</b>	15/Nedd4
<b>Immunogen:</b>	Mouse Nedd4 aa. 173-262
<b>Isotype:</b>	Mouse IgG1
<b>Reactivity:</b>	QC Testing: Mouse
<b>Target MW:</b>	110 kDa
<b>Storage Buffer:</b>	Aqueous buffered solution containing BSA, glycerol, and ≤0.09% sodium azide.


**Description**

Regulated protein degradation involves conjugation of ubiquitin to proteins for degradation by the 26S proteasome. Protein ubiquitination is the end product of a multi-enzyme cascade that involves sequential thiol-ester bond interactions between ubiquitin and enzymes termed E1-E3. Neuronal precursor cell-expressed developmentally down-regulated 4 (Nedd4) is a multimodular ubiquitin protein ligase E3 that contains a Ca<sup>2+</sup>-lipid binding domain (CaLB), 3 WW domains, and a C-terminal ubiquitin protein ligase hect domain. The CaLB is characteristic of C2 domain-containing proteins, such as PKC, phospholipase C, phospholipase A2, and synaptogamin. Nedd4 is expressed differentially in the fetal and adult brain and its highest expression is in lung and kidney. Although it localizes in the cytoplasm, increased intracellular Ca<sup>2+</sup> induces plasma membrane association of Nedd4. In addition, increases in intracellular Na<sup>+</sup> leads to Nedd4 binding to epithelial Na<sup>+</sup> channels via its WW domains, which results in targeting of Na<sup>+</sup> channels for degradation. Thus, the subcellular localization and protein interactions of Nedd4 are regulated in a Ca<sup>2+</sup>- and Na<sup>+</sup>-dependent manner.

This antibody is routinely tested by western blot analysis. Other applications were tested at BD Biosciences Pharmingen during antibody development only or reported in the literature.



**Western blot analysis of Nedd4 on a BC3H1 cell lysate (murine skeletal muscle cells). Lane 1: 1:1000, lane 2: 1:2000, lane 3: 1:4000 dilution of the anti-Nedd4 antibody.**



**Immunofluorescence staining of RSV-3T3 cells.**

**Preparation and Storage**

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

Store undiluted at -20° C.

**BD Biosciences**

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country-specific contact information, visit [bdbiosciences.com/how\\_to\\_order/](http://bdbiosciences.com/how_to_order/)

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2006 BD



**BD**

**BD Biosciences**

## Application Notes

### Application

Western blot	Routinely Tested
Immunofluorescence	Tested During Development

## Suggested Companion Products

<u>Catalog Number</u>	<u>Name</u>	<u>Size</u>	<u>Clone</u>
554002	HRP Goat Anti-Mouse Igs	1.0 ml	(none)
554001	FITC Goat Anti-Mouse Igs	0.5 mg	Polyclonal

## Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to [www.bdbiosciences.com/pharming/en/protocols](http://www.bdbiosciences.com/pharming/en/protocols) for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
4. Source of all serum proteins is from USDA inspected abattoirs located in the United States.

## References

Harvey KF, Dinudom A, Komwatana P, et al. All three WW domains of murine Nedd4 are involved in the regulation of epithelial sodium channels by intracellular Na<sup>+</sup>. *J Biol Chem.* 1999; 274(18):12525-12530.(Biology)

Hatakeyama S, Jensen JP, Weissman AM. Subcellular localization and ubiquitin-conjugating enzyme (E2) interactions of mammalian HECT family ubiquitin protein ligases. *J Biol Chem.* 1997; 272(24):15085-15092.(Biology)

Peruzzi F, Prisco M, Morrione A, Valentini B, Baserga R. Anti-apoptotic signaling of the insulin-like growth factor-I receptor through mitochondrial translocation of c-Raf and Nedd4. *J Biol Chem.* 2001; 276(28):25990-25996.(Biology: Western blot)

Plant PJ, Yeger H, Staub O, Howard P, Rotin D. The C2 domain of the ubiquitin protein ligase Nedd4 mediates Ca<sup>2+</sup>-dependent plasma membrane localization. *J Biol Chem.* 1997; 272(51):32329-32336.(Biology)