


Technical Data Sheet

Purified Mouse Anti-Stat 6 (pY641)**Product Information**

| | |
|-------------------------|--|
| Material Number: | 611567 |
| Size: | 150 µg |
| Concentration: | 250 µg/ml |
| Clone: | 18/P-Stat6 |
| Immunogen: | Phosphorylated Human Stat6 (pY641) |
| Isotype: | Mouse IgG2a |
| Reactivity: | QC Testing: Human |
| Target MW: | 100 kDa |
| Storage Buffer: | Aqueous buffered solution containing BSA, glycerol, and ≤0.09% sodium azide. |

Description

Interleukin-4 (IL-4), a major immunoregulatory cytokine, is secreted by activated T lymphocytes, basophils, and mast cells and plays an important role in modulating T helper cell lineage development. It induces specific gene expression via the tyrosine phosphorylation of Stat6 at tyrosine 641 (Y641). Stat6, a member of the signal transducers and activators of transcription protein family, mediates signals for IL-4 and, possibly, IL-13. While Stat6 is widely expressed in human tissues, it exhibits elevated expression in peripheral blood lymphocytes, colon, intestine, ovary, prostate, thymus, spleen, kidney, liver, lung, and placenta. Following cytokine receptor ligation, Jak kinases are activated and phosphorylate the cytoplasmic tails of the oligomerized receptors. The SH3:SH2 domain of Stat6 associates with tyrosine-phosphorylated IL-4 receptor and the proximal Jak kinase phosphorylates Stat6 at Y641 on the C-terminal side of the SH2 domain. Stat6 is then released from the receptor, dimerizes, and is thought to contact the basal transcription machinery by binding to p300/CBP. Thus, Stat6 mediates the IL-4 signal and is essential for the proper development of adaptive immunity.



Immunofluorescent labelling of phosphorylated Stat6 in endothelial cells exposed to 10 ng/ml IL-4 for 1 hour.



Human Endothelial cells were either untreated (left column) or treated with (right column) IL-4 (10 ng/ml) for 1 hour at 37°C. The top panel was probed with Stat6 (Cat. No. 611290) and the bottom panel was probed with Stat6 (pY641) (Cat. No. 611820).

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. Store undiluted at -20°C.

BD Biosciences

bdbiosciences.com

| | | | | | |
|---------------|--------------|---------------|--------------|--------------|-------------------------|
| United States | Canada | Europe | Japan | Asia Pacific | Latin America/Caribbean |
| 877.232.8995 | 888.259.0187 | 32.53.720.550 | 0120.8555.90 | 65.6861.0633 | 55.11.5185.9995 |

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2008 BD



Application Notes

Application

| | |
|--------------------|---------------------------|
| Western blot | Routinely Tested |
| Immunofluorescence | Tested During Development |

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to www.bdbiosciences.com/pharming/en/protocols for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
4. Source of all serum proteins is from USDA inspected abattoirs located in the United States.

References

- Hou J, Schindler U, Henzel WJ, Ho TC, Brasseur M, McKnight SL. An interleukin-4-induced transcription factor: IL-4 Stat. *Science*. 1994; 265(5179):1701-1706. (Biology)
- Mikita T, Campbell D, Wu P, Williamson K, Schindler U. Requirements for interleukin-4-induced gene expression and functional characterization of Stat6. *Mol Cell Biol*. 1996; 16(10):5811-5820.(Biology)
- Quelle FW, Shimoda K, Thierfelder W, et al.. Cloning of murine Stat6 and human Stat6. Stat proteins that are tyrosine phosphorylated in responses to IL-4 and IL-3 but are not required for mitogenesis. *Mol Cell Biol*. 1995; 15(6):3336-3343.(Biology)