

Technical Data Sheet


Purified Mouse Anti-Serine Racemase**Product Information**

Material Number:	612053
Size:	150 µg
Concentration:	250 µg/ml
Clone:	29/Serine Racemase
Immunogen:	Mouse serine racemase aa. 127-248
Isotype:	Mouse IgG1
Reactivity:	QC Testing: Mouse Tested in Development: Rat
Target MW:	38 kDa
Storage Buffer:	Aqueous buffered solution containing BSA, glycerol, and ≤0.09% sodium azide.

Description

The majority of synapses in the central nervous system utilize glutamate as a neurotransmitter to produce rapid neuronal excitation. Glutamate has a diverse array of receptors that can be categorized into two groups: ionotropic and metabotropic. The ionotropic receptors are subdivided into two distinct types: 1) receptors for N-methyl D-aspartate (NMDAR) and 2) non-NMDA receptors for AMPA and kainate. NMDA receptors require coactivation at both glutamate and glycine sites. D-serine is a D-amino acid found in mammalian tissues that can act as a potent ligand for the glycine site on NMDA receptors. D-serine is made by the enzyme serine racemase, which is a member of the pyridoxal-5' phosphate (PLP)-dependent enzyme family. Serine racemase mRNA is expressed in brain and liver, and serine racemase protein is expressed in glial cells. Degradation of D-serine by D-amino acid oxidase attenuates NMDA receptor-mediated calcium influx, and implicates D-serine as an endogenous modulator of NMDA receptor function. Thus, glial cell production of D-serine via serine racemase activity may be an important system for modulation of synaptic transmission.

This antibody is routinely tested by western blot analysis. Other applications were tested at BD Biosciences Pharmingen during antibody development only or reported in the literature.



Western blot analysis of serine racemase in a mouse cerebrum lysate. Lane 1: 1:500, lane 2: 1:1000, lane 3: 1:2000 dilution of the anti-serine racemase antibody.

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. Store undiluted at -20° C.

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2006 BD

**BD****BD Biosciences**

Application Notes

Application

Western blot	Routinely Tested
Immunofluorescence	Not Recommended

Suggested Companion Products

Catalog Number	Name	Size	Clone
611455	Mouse Cerebrum Lysate	500 µg	(none)
554002	HRP Goat Anti-Mouse Igs	1.0 ml	(none)

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to www.bdbiosciences.com/pharming/en/protocols for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
4. Source of all serum proteins is from USDA inspected abattoirs located in the United States.

References

Mothet JP, Parent AT, Wolosker H, et al. D-serine is an endogenous ligand for the glycine site of the N-methyl-D-aspartate receptor. *Proc Natl Acad Sci U S A*. 2000; 97(9):4926-4931.(Biology)
Wolosker H, Blackshaw S, Snyder SH. Serine racemase: a glial enzyme synthesizing D-serine to regulate glutamate-N-methyl-D-aspartate neurotransmission. *Proc Natl Acad Sci U S A*. 1999; 96(23):13409-13414.(Biology)