Technical Data Sheet **Purified Mouse Anti-** α-Methylacyl-CoA Racemase

Product Information			
Material Number:	612082		
Alternate Name:	AMACR		
Size:	50 µg		
Concentration:	250 μg/ml		
Clone:	15/α-Methylacyl-CoA Racemase		
Immunogen:	Human α-Methylacyl-CoA Racemase aa.139-229		
Isotype:	Mouse IgG1		
Reactivity:	QC Testing: Mouse Tested in Development: Human, Rat		
Target MW:	42 kDa		
Storage Buffer:	Aqueous buffered solution containing BSA, glycerol, and $\leq 0.09\%$ sodium azide.		

Description

 α -methylacyl-CoA Racemase (AMACR) catalyzes the racemization of α -methyl-branched carboxylic acid coenzyme A thioesters. This enzymatic function is involved with biosynthesis of bile acids in the mitochondria and peroxisomes. AMACR converts pristanoyl-CoA and C27-bile acyl-CoAs to their (S)-stereoisomers, which are the only stereoisomers degraded by peroxisomal α -oxidation. Interestingly, AMACR mutations have been linked to some sensory motor neuropathies where accumulation of fatty acids and AMACR deficiencies correlate with pathogenesis. AMACR contains an N-terminal region required for mitochondrial localization, and a C-terminal peroxisomal targeting signal type 1 (PTS). AMACR mRNA is expressed preferentially in human, rat, and mouse liver and kidney, but the percentage of AMACR enzymatic activity found in the mitochondria relative to the peroxisomes differs depending on the species. Thus, AMACR is an enzyme critical for fatty acid degradation, and bile formation.



Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. Store undiluted at -20°C.

Application Notes

Application

Western blot	Routinely Tested
Immunofluorescence	Not Recommended

BD Biosciences

bdbiosciences.com							
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995		
For country-spe	ecific contact in	formation, visit	bdbiosciences.co	m/how_to_orde	r/		
Conditions: The ir of any patents. Bl use of our produc product or as a co written authoriza For Research Use	nformation disclose D Biosciences will n cts. Purchase does n imponent of anoth tion of Becton Dick Only. Not for use in	d herein is not to b ot be held responsi not include or carry er product. Any us kinson and Compan n diagnostic or there	e construed as a rec ble for patent infrin any right to resell or e of this product oth y is strictly prohibite apeutic procedures.	ommendation to us gement or other vic r transfer this produ her than the permitt ed. Not for resale.	e the above product in violation vlations that may occur with the ct either as a stand-alone red use without the express		
BD, BD Logo and	all other trademar	ks are the property	of Becton, Dickinsor	n and Company. ©20	08 BD		



Recommended Assay Procedure:

Western blot: Please refer to http://www.bdbiosciences.com/pharmingen/protocols/Western_Blotting.shtml

Suggested Companion Products

Catalog Number	Name	Size	Clone
611458	Mouse Liver Lysate	500 μg	(none)
554002	HRP Goat Anti-Mouse Ig	1.0 ml	(none)

Product Notices

- 1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
- 2. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
- 3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
- 4. Source of all serum proteins is from USDA inspected abattoirs located in the United States.

References

Ferdinandusse S, Denis S, Clayton PT. Mutations in the gene encoding peroxisomal alpha-methylacyl-CoA racemase cause adult-onset sensory motor neuropathy. Nat Genet. 2000; 24(2):188-191.(Biology)

Kotti TJ, Savolainen K, Helander HM. In mouse alpha -methylacyl-CoA racemase, the same gene product is simultaneously located in mitochondria and peroxisomes. J Biol Chem. 2000; 275(27):20887-20895.(Biology)