

Technical Data Sheet

Purified Mouse Anti-Stat1 (pY701)**Product Information**

Material Number:	612233
Size:	150 µg
Concentration:	250 µg/ml
Clone:	4a
Immunogen:	Phosphorylated Human Stat1 Peptide
Isotype:	Mouse IgG2a
Reactivity:	QC Testing: Human Tested in Development: Mouse
Target MW:	84/91 kDa
Storage Buffer:	Aqueous buffered solution containing BSA, glycerol, and ≤0.09% sodium azide.


Description

Stat (Signal transducer and activators of transcription) proteins are critical mediators of the biologic activity of cytokines, including interleukins, interferons, erythropoietin, and growth factors. Ligand-receptor interaction leads to activation of constitutively associated JAK family kinases and subsequent recruitment/activation of Stat proteins by tyrosine phosphorylation. Active Stat proteins then move to the nucleus to promote transcription of cytokine-inducible genes. Seven Stat proteins have been cloned, each of which is differentially expressed and/or activated in a cytokine-specific and cell type-specific manner. Stat1 and Stat2 are components of the ISGF3 (Interferon-Stimulated Gene Factor 3) complex, which is the primary transcription activator induced by the binding of the interferon to a specific cell-surface receptor. Stat1 has two alternatively spliced isoforms, 91-kDa Stat1 α and 84-kDa Stat1 β ; Stat1 α has 38 additional C-terminal amino acids. In response to the binding of IFN α , IFN γ , EGF, PDGF, or CSF-1 to their respective receptors, the Stat1 subunits become tyrosine-phosphorylated at Y701, and the complex is translocated to the nucleus. This results in the formation of an active complex that includes the DNA-binding p48 subunit. This complex is responsible for modulating the transcription of the interferon-stimulated genes (ISGs).

The 4a monoclonal antibody recognizes the phosphorylated Y701 in Stat1 α and Stat1 β .



A431 cells were either left untreated (left lane) or treated (right lane) with 100 ng/ml EGF for 5 minutes at 37°C. The top panel was probed with Stat1 (Cat. No. 610115) and the bottom was probed with Stat1 (pY701) (Cat. No. 612233).



A431 cells were serum starved and treated with EGF (100 ng/ml) for 5 minutes, then fixed in 3.75% paraformaldehyde with 0.2% Triton X-100. Immunofluorescent staining was performed with Stat1 (Cat. No. 610115) and Stat1 (pY701) (Cat. No. 612233).

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

Store undiluted at -20°C.

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2008 BD



Application Notes

Application

Western blot	Routinely Tested
Immunofluorescence	Tested During Development
Immunoprecipitation	Tested During Development

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
4. Source of all serum proteins is from USDA inspected abattoirs located in the United States.

References

Darnell JE Jr. STATs and gene regulation. *Science*. 1997; 277(5332):1630-1635.(Biology)
Fu XY, Zhang JJ. Transcription factor p91 interacts with the epidermal growth factor receptor and mediates activation of the c-fos gene promoter. *Cell*. 1993; 74(6):1135-1145.(Biology)