


## Technical Data Sheet

**Purified Mouse Anti-p38 MAPK (pT180/pY182)****Product Information**

<b>Material Number:</b>	<b>612281</b>
<b>Size:</b>	150 µg
<b>Concentration:</b>	250 µg/ml
<b>Clone:</b>	30/p38 MAPK (pT180/pY182)
<b>Immunogen:</b>	Human p38 MAPK (pT180/pY182)
<b>Isotype:</b>	Mouse IgG1
<b>Reactivity:</b>	QC Testing: Human Tested in Development: Mouse, Rat
<b>Target MW:</b>	38-42 kDa
<b>Storage Buffer:</b>	Aqueous buffered solution containing BSA, glycerol, and ≤0.09% sodium azide.

**Description**

Activation of the immune and inflammatory responses often involves the recognition of bacterial endotoxin (lipopolysaccharide or LPS). Binding of LPS by monocytic cells results in the production and release of proinflammatory cytokines, such as IL-1 and TNF- $\alpha$ . LPS-induced signaling cascades involve members of the Ser/Thr protein kinase family known as the mitogen activated protein kinases (MAPKs). MAPK signal transduction pathways mediate the effects of various extracellular stimuli on biological processes such as proliferation, differentiation, and death. The p38 MAP kinases include p38 $\alpha$ ,  $\beta$ ,  $\gamma$ , and  $\delta$ . These ser/thr kinases are activated by dual phosphorylation on Thr and Tyr within the motif Thr-Gly-Tyr located in kinase subdomain VIII. Activation of p38 MAPK is mediated specifically by the MAP kinase kinases, MKK3, MKK4, and MKK6. This leads to the activation of multiple transcription factors (NF- $\kappa$ B, ATF-2, Elk-1, and CHOP) that induce expression of many different genes, including proinflammatory cytokine genes. Thus, p38 MAPKs are central kinases in multiple signal transduction pathways.



**Western blot analysis for p38 MAPK (pT180/pY182) (left figure).** HeLa cells (Human cervical epitheloid carcinoma; ATCC CCL-2) were either left untreated (lane 1) or treated with 25 µg/mL anisomycin, an antibiotic and protein synthesis inhibitor, for 15 min at 37°C (lane 2). The top panel was probed with a mouse anti-p38 $\alpha$  antibody (cat.# 612168) and the bottom was probed with the mouse anti-p38 (pT180/pY182) antibody.

**Immunofluorescence staining for p38 MAPK (pT180/pY182) (right figure).** HeLa cells (Human cervical epitheloid carcinoma; ATCC CCL-2) were either untreated (upper left and bottom left panels) or were treated for 15 min with 25 µg/mL anisomycin, an antibiotic and protein synthesis inhibitor (upper right and bottom right panels). Cells were fixed in 3.75% paraformaldehyde and permeabilized with 0.2% Triton-X 100. Immunofluorescent staining was performed with either a mouse anti-p38 $\alpha$  antibody (cat.#612168) (upper left and upper right panels) or with the mouse anti-p38 MAPK (pT180/pY182) antibody (bottom left and bottom right panels).

**BD Biosciences**

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country-specific contact information, visit [bdbiosciences.com/how\\_to\\_order/](http://bdbiosciences.com/how_to_order/)

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2008 BD



## Preparation and Storage

Store undiluted at -20°C.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

## Application Notes

### Application

Western blot	Routinely Tested
Flow cytometry	Tested During Development
Immunofluorescence	Tested During Development

### Recommended Assay Procedure:

**Western blot:** Please refer to [http://www.bdbiosciences.com/pharmingen/protocols/Western\\_Blotting.shtml](http://www.bdbiosciences.com/pharmingen/protocols/Western_Blotting.shtml)

## Suggested Companion Products

Catalog Number	Name	Size	Clone
611449	HeLa Cell Lysate	500 µg	(none)
611692	HeLa + Anisomycin Cell Lysate	500 µg	(none)
554002	HRP Goat Anti-Mouse Ig	1.0 ml	(none)
554001	FITC Goat Anti-Mouse Ig	0.5 mg	Polyclonal
612168	Purified Mouse Anti-p38α	50 µg	27/p38α/SAPK2a

## Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Source of all serum proteins is from USDA inspected abattoirs located in the United States.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
4. Please refer to [www.bdbiosciences.com/pharmingen/protocols](http://www.bdbiosciences.com/pharmingen/protocols) for technical protocols.

## References

Brunet A, Pouyssegur J. Identification of MAP kinase domains by redirecting stress signals into growth factor responses. *Science*. 1996; 272(5268):1652-1655. (Biology)

Han J, Lee JD, Bibbs L, Ulevitch RJ. A MAP kinase targeted by endotoxin and hyperosmolarity in mammalian cells. *Science*. 1994; 265(5173):808-811. (Biology)

Winston BW, Chan ED, Johnson GL, Riches DW. Activation of p38mapk, MKK3, and MKK4 by TNF-alpha in mouse bone marrow-derived macrophages. *J Immunol*. 1997; 159(9):4491-4497. (Biology)