Technical Data Sheet

Purified Mouse Anti-Hsp75/TRAP1

Product Information

Material Number: 612344 Alternate Name: TRAP1 Size 50 μg **Concentration:** $250 \mu g/ml$ Clone: 42/Hsp75

Human Hsp75/TRAP1 aa. 253-464 Immunogen:

Isotype: Mouse IgG1 Reactivity: QC Testing: Human


Tested in Development: Dog, Rat, Mouse


Target MW:

Storage Buffer: Aqueous buffered solution containing BSA, glycerol, and ≤0.09% sodium

Description

The hsp90 family of molecular chaperones is involved in a number of signal transduction pathways, including cell division, protein folding and transport. A new member of this family was recently identified as TRAP1 (Tumor necrosis factor receptor-associated protein) or hsp75. TRAP1 is a mitochondrial protein that is an ATPase and ATP-binding protein, rendering it similar to hsp90. There is a high degree of homology between TRAP1 and hsp90 in 6 domains, however one major difference is that a highly charged domain present in hsp90 is absent in TRAP1 TRAP1 has been shown to interact with the intracellular domain of the type 1 receptor for tumor necrosis factor (TNFR-1IC) and to the retinoblastoma protein (Rb). Signaling through TNFR-1 induces antiviral responses, fibroblast proliferation, cytotoxicity, etc. TRAP1 binds TNFR-1IC outside of the death domain leading to disruption in the signaling by TNFR-1. Rb has been shown to play a pivotal role in cellular differentiation and progression through the cell cycle. TRAP1 associates with Rb via an LxCxE motif, which is common among proteins that interact with Rb through SV40 T-antigen binding domain. Thus, TRAP1 is a new member of the hsp90 family that interacts with Rb and TNFR-1IC.





Western blot analysis of Hsp75 on HeLa lysate. Lane 1: 1:500, lane 2: 1:1000, lane 3: 1:2000 dilution of Hsp75.

Immunofluorescence staining of HeLa cells.

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. Store undiluted at -20°C.

Application Notes

Application

| Western blot | Routinely Tested | |
|--------------------|---------------------------|--|
| Immunofluorescence | Tested During Development | |

BD Biosciences

bdbiosciences.com

United States Asia Pacific Latin America/Caribbean Canada Europe 877.232.8995 888.259.0187 32.53.720.550 0120.8555.90 65.6861.0633 55.11.5185.9995

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2008 BD



Suggested Companion Products

| Catalog Number | Name Name | Size | Clone | |
|----------------|-------------------------|--------|------------|--|
| 554002 | HRP Goat Anti-Mouse Ig | 1.0 ml | (none) | |
| 554001 | FITC Goat Anti-Mouse Ig | 0.5 mg | Polyclonal | |
| 611449 | HeLa Cell Lysate | 500 μg | (none) | |

Product Notices

- 1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
- 2. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
- Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
- 4. Source of all serum proteins is from USDA inspected abattoirs located in the United States.

References

Chen CF, Chen Y, Dai K, Chen PL, Riley DJ, Lee WH. A new member of the hsp90 family of molecular chaperones interacts with the retinoblastoma protein during mitosis and after heat shock. *Mol Cell Biol.* 1996; 16(9):4691-4699.(Biology)

Felts SJ, Owen BA, Nguyen P, Trepel J, Donner DB, Toft DO. The hsp90-related protein TRAP1 is a mitochondrial protein with distinct functional properties. *J Biol Chem.* 2000; 275(5):3305-3312.(Biology)

Song HY, Dunbar JD, Zhang YX, Guo D, Donner DB. Identification of a protein with homology to hsp90 that binds the type 1 tumor necrosis factor receptor. *J Biol Chem.* 1995; 270(8):3574-3581.(Biology)

612344 Rev. 1 Page 2 of 2