


## Technical Data Sheet

**Purified Mouse Anti-Lck (pY505)****Product Information**

|                         |  |
|-------------------------|--|
| <b>Material Number:</b> | 612390   |
| <b>Size:</b>            | 50 µg  |
| <b>Concentration:</b>   | 250 µg/ml  |
| <b>Clone:</b>           | 4/Lck (pY505)  |
| <b>Immunogen:</b>       | Human Lck (pY505)  |
| <b>Isotype:</b>         | Mouse IgG1   |
| <b>Reactivity:</b>      | QC Testing: Human<br>Tested in Development: Rat, Mouse                       |
| <b>Target MW:</b>       | 56 kDa   |
| <b>Storage Buffer:</b>  | Aqueous buffered solution containing BSA, glycerol, and ≤0.09% sodium azide. |

**Description**

Protein tyrosine phosphorylation is an essential step in the signal transduction cascade leading to T cell antigen receptor (TCR) activation. Lck is a protein kinase and a member of the src family of cytoplasmic protein-tyrosine kinases (PTKs). Members of this family have several common features: 1) unique N-terminal domains, 2) attachment to cellular membranes through a myristylated N-terminus, and 3) homologous SH2, SH3, and catalytic domains. The unique N-terminal domain of Lck interacts with the cytoplasmic tails of the CD4 and CD8 cell surface glycoproteins. CD4 and CD8 bind to surface MHC class II and class I molecules, respectively. Lck is regulated by both kinases and phosphatases. Autophosphorylation at Y394 leads to conformational changes in the catalytic domain, which induces kinase activity. Repression of Lck occurs via phosphorylation at Y505, located near the carboxy-terminus. Phosphorylation of this tyrosine site is mediated by the Csk family of PTKs. Upon phosphorylation at this site, Lck associates with the SH2 domain in the amino-terminus, thus keeping the protein biologically inactive. Lck activity and regulation is critical for activation and development of T cells.



*Jurkat cells were either untreated (lane 1) or treated (lane 2) with Anti-CD3 for 15 minutes at 37°C. The top panel was probed with Lck (cat. #610097) and the bottom panel was probed with Lck (pY505) (cat. #612390).*

**Preparation and Storage**

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. Store undiluted at -20° C.

**Application Notes****Application**

|                |                           |
|----------------|---------------------------|
| Western blot   | Routinely Tested          |
| Flow cytometry | Tested During Development |

**BD Biosciences**

[www.bdbiosciences.com](http://www.bdbiosciences.com)

United States 877.232.8995 Canada 888.259.0187 Europe 32.53.720.550 Japan 0120.8555.90 Asia Pacific 65.6861.0633 Latin America/Caribbean 55.11.5185.9995

For country-specific contact information, visit [www.bdbiosciences.com/how\\_to\\_order/](http://www.bdbiosciences.com/how_to_order/)

*Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.*

*For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.*

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2007 BD



## Suggested Companion Products

| <u>Catalog Number</u> | <u>Name</u>                 | <u>Size</u> | <u>Clone</u> |
|-----------------------|-----------------------------|-------------|--------------|
| 610097                | Purified Mouse Anti-Lck     | 50 µg       | 28/Lck       |
| 611755                | Jurkat + Pervanadate Lysate | 500 µg      | (none)       |

## Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to [www.bdbiosciences.com/pharmingen/protocols](http://www.bdbiosciences.com/pharmingen/protocols) for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
4. Source of all serum proteins is from USDA inspected abattoirs located in the United States.

## References

Hardwick JS, Sefton BM. The activated form of the Lck tyrosine protein kinase in cells exposed to hydrogen peroxide is phosphorylated at both Tyr-394 and Tyr-505. *J Biol Chem.* 1997; 272(41):25429-25432.(Biology)

Lee-Fruman KK, Collins TL, Burakoff SJ. Role of the Lck Src homology 2 and 3 domains in protein tyrosine phosphorylation. *J Biol Chem.* 1996; 271(40):25003-25012.(Biology)

Wang B, Lemay S, Tsai S, Veillette A. SH2 domain-mediated interaction of inhibitory protein tyrosine kinase Csk with protein tyrosine phosphatase-HSCF. *Mol Cell Biol.* 2001; 21(4):1077-1088.(Biology)