

**TABLE OF CONTENTS**

<b>1.0</b>	<b>Overview of GeneBLAzer<sup>®</sup> Technology</b> .....	<b>1</b>
<b>2.0</b>	<b>Materials Supplied</b> .....	<b>1</b>
<b>3.0</b>	<b>Materials Required, but Not Supplied</b> .....	<b>2</b>
3.1	Optional Equipment and Materials .....	2
<b>4.0</b>	<b>Cell Culture Conditions</b> .....	<b>3</b>
4.1	Media Required .....	3
4.2	Growth Conditions .....	3
<b>5.0</b>	<b>Assay Procedure</b> .....	<b>4</b>
5.1	Quick Reference Guide .....	4
5.2	Detailed Assay Protocol .....	4
5.3	Detection .....	5
<b>6.0</b>	<b>Data Analysis</b> .....	<b>6</b>
6.1	Background Subtraction .....	6
6.2	Visual Observation of Intracellular Beta-lactamase Activity Using LiveBLAzer <sup>™</sup> -FRET B/G Substrate (CCF4-AM).....	6
6.3	Representative Data .....	6
<b>7.0</b>	<b>Detailed Cell Handling Procedures</b> .....	<b>7</b>
7.1	Thawing Method .....	7
7.2	Propagation Method .....	7
7.3	Freezing Method.....	7
<b>8.0</b>	<b>References</b> .....	<b>7</b>
<b>9.0</b>	<b>Purchaser notification</b> .....	<b>8</b>

**1.0 OVERVIEW OF GENEBLAZER<sup>®</sup> TECHNOLOGY**

GeneBLAzer<sup>®</sup> Beta-lactamase Reporter Technology provides a highly accurate, sensitive, and easy to use method of monitoring cellular responses to drug candidates or other stimuli (1). The core of the GeneBLAzer<sup>®</sup> Technology is a Fluorescence Resonance Energy Transfer (FRET) substrate that generates a ratiometric reporter response with minimal experimental noise. In addition to the two-color (blue/green) readout of stimulated and unstimulated cells, this ratiometric method reduces the absolute and relative errors that can mask the underlying biological response of interest. Such errors include variations in cell number, transfection efficiency, substrate concentration, excitation path length, fluorescence detectors, and volume changes. The GeneBLAzer<sup>®</sup> Beta-lactamase Reporter Technology has been proven effective in high-throughput screening (HTS) campaigns for a range of target classes, including G-protein coupled receptors (GPCRs) (2, 3), nuclear receptors (4-6), and kinase signaling pathways (7).

**2.0 MATERIALS SUPPLIED**

<b>Cell Line Name:</b>	AP1- <i>bla</i> A375
<b>Description:</b>	CellSensor <sup>®</sup> AP1- <i>bla</i> A375 cells contain a beta-lactamase reporter gene under control of the AP1 Response Element that has been stably integrated into A375 cells. AP1- <i>bla</i> A375 cells constitutively express beta-lactamase, which can be inhibited with Raf1 kinase inhibitor.
<b>Product Number:</b>	K1287
<b>Shipping Condition:</b>	Dry Ice
<b>Storage Condition:</b>	Liquid nitrogen. <b>Immediately upon receipt, cells must be stored in liquid nitrogen or thawed for immediate use. Cells stored at -80°C can quickly lose viability.</b>
<b>Quantity:</b>	~2,000,000 (2 × 10 <sup>6</sup> cells/ml)
<b>Application:</b>	Detection of inhibitors for B-Raf wt and V600E signaling pathway.
<b>Growth Properties:</b>	Adherent
<b>Cell Phenotype:</b>	Melanoma
<b>Selection Marker:</b>	Blasticidin (5 µg/ml)
<b>Vector Used:</b>	pLenti- <i>bsd</i> /AP1- <i>bla</i> Vector
<b>Mycoplasma Testing:</b>	Negative
<b>BioSafety Level:</b>	1

### 3.0 MATERIALS REQUIRED, BUT NOT SUPPLIED

Media/Reagents	Recommended Source	Part #
LiveBLAzer™-FRET B/G Loading Kit, containing: LiveBLAzer™-FRET B/G Substrate (CCF4-AM), DMSO, Solution B, and Solution C	Invitrogen	K1095 (0.2 mg) K1096 (1 mg) K1030 (5 mg)
Recovery™ Cell Culture Freezing Medium	Invitrogen	12648-010
DMEM + GlutaMax™-1	Invitrogen	10569-010
Opti-MEM® Reduced Serum Medium	Invitrogen	11058-021
DMSO	Fluka	41647
Fetal Bovine Serum (FBS), dialyzed, tissue-culture grade ( <b>DO NOT SUBSTITUTE!</b> )	Invitrogen	26400-044
Nonessential amino acids (NEAA)	Invitrogen	11140-050
Penicillin/Streptomycin (antibiotic)	Invitrogen	15140-122
Sodium Pyruvate	Invitrogen	11360-070
Raf1 Inhibitor I	EMD	553008
0.05% Trypsin/EDTA	Invitrogen	25300-054
Solution D	Invitrogen	K1156
Blasticidin (antibiotic)	Invitrogen	R210-01

Consumables	Recommended Source	Part #
Black-wall, clear-bottom, 384-well assay plates (with low fluorescence background)	Corning Life Sciences	3712
Compressed air	Various	—

Equipment	Recommended Source
Fluorescence plate reader with bottom-read capability	Various
Filters, if required for plate reader (see <b>Section 5.3</b> )	Chroma Technology Corp.

**Note:** If you do not have access to a fluorescence plate reader with bottom-read capabilities, contact our Technical Support for options of other beta-lactamase substrates that can be read with top-reading instruments.

#### 3.1 Optional Equipment and Materials

- Epifluorescence- or fluorescence-equipped microscope with appropriate filters
- Microplate centrifuge

## 4.0 CELL CULTURE CONDITIONS

### 4.1 Media Required

Component	Thaw Medium	Growth Medium	Assay Medium	Freezing Medium
DMEM + GlutaMax™-1	500 ml bottle	500 ml bottle	—	—
Opti-MEM Medium	—	—	500 ml bottle	—
Dialyzed FBS	50 ml	50 ml	2.5 ml	—
NEAA (10 mM)	—	—	5 ml	—
Sodium Pyruvate (100 mM)	—	—	5 ml	—
Penicillin (10,000 U/ml) / Streptomycin (10,000 µg/ml)	5 ml	5 ml	5 ml	—
Blasticidin	—	5 µg/ml	—	—
Recovery™ Cell Culture Freezing Medium		—	—	100%

**Note:** We prepare our media by adding the listed components directly to the medium bottle. Blasticidin can be added directly to the cell culture flask to reach 5 µg/ml. Similar methods are suitable.

**Note:** Unless otherwise stated, have all media and solutions at least at room temperature (we recommend 37°C for optimal performance) before adding them to the cells.

### 4.2 Growth Conditions

For detailed cell growth and maintenance directions, see **Section 7.0**.

**Note:** We recommend passing cells for three passages after thawing before using them in the beta-lactamase assay.

1. Thaw cells in Thaw Medium **without Blasticidin** and culture them in Growth Medium with Blasticidin. Pass or feed cells at least three times a week and maintain them in a 37°C/5% CO<sub>2</sub> incubator. Maintain cells between 10% to 90% confluency. Do not allow cells to reach confluence.
2. Freeze cells at  $2 \times 10^6$  cells/ml in Recovery™ Cell Culture Freezing Medium.

## 5.0 ASSAY PROCEDURE

The following instructions outline the recommended procedure for monitoring B-RAF V600E induced MAPK signaling using LiveBLAzer™-FRET B/G Substrate as the readout. If you use alternative substrates (e.g., ToxBLAzer™ DualScreen, or LyticBLAzer™ Loading kits), follow the loading protocol provided with the product.

### Note:

- We recommend using 384-well, black-wall, clear-bottom assay plates with low fluorescence background.
- Some solvents may affect assay performance. Assess the effect of a test compound solvent before screening. The cell stimulation described below is carried out in the presence of 0.5% DMSO to simulate the effect that a test compound solvent might have on the assay. If you use other solvents and/or solvent concentrations, change the following assay conditions and optimize appropriately.

### 5.1 Quick Reference Guide

For more detailed protocol information, see **Section 5.2**. Plate layouts and experimental outlines will vary; in screening mode, we recommend using at least three wells for each condition: Unstimulated, Stimulated, and Cell-free.

	Untreated Wells	Compound-treated Wells	Cell-free wells
<b>Step 1</b> Plate cells	32 $\mu$ l cells suspended in Assay Medium (10,000 cells/well)	32 $\mu$ l cells suspended in Assay Medium (10,000 cells/well)	32 $\mu$ l Assay Medium (no cells)
<b>Step 2</b> Add DMSO	Add 4 $\mu$ l of 4% DMSO in Assay Medium to each well		
<b>Step 3</b> Add compound	4 $\mu$ l 1% DMSO in Assay Medium to each well	4 $\mu$ l 10X compound dilution in Assay Medium to each well	4 $\mu$ l 1% DMSO in Assay Medium to each well
<b>Step 4</b> Incubate cells	Incubate the plate at 37°C/5% CO <sub>2</sub> for 17 hours.		
<b>Step 5</b> Prepare 6X Substrate Mixture	6 $\mu$ l 1 mM LiveBLAzer™-FRET B/G Substrate (CCF4-AM) in dry DMSO + 60 $\mu$ l Solution B, mix. Add 874 $\mu$ l Solution C and 60 $\mu$ l Solution D, mix.		
<b>Step 6</b> Load Substrate Mixture	8 $\mu$ l per well		
<b>Step 7</b> Incubate Substrate + cells	2.5 hours at room temperature in the dark.		
<b>Step 8</b> Detect Activity	See <b>Section 5.3</b>		
<b>Step 9</b> Analyze data	See <b>Section 6.0</b>		

### 5.2 Detailed Assay Protocol

Plate layouts and experimental outlines will vary; in screening mode, we recommend using at least three wells for each control: Untreated Control, Compound-treated Control, and Cell-free Control.

#### 5.2.1 Precautions

- Work on a dust-free, clean surface. Always handle the 384-well, black-wall, clear-bottom assay plate by the sides; do not touch the clear bottom of the assay plate.
- If pipetting manually, you may need to centrifuge the plate briefly at room temperature (30 seconds at 14  $\times$  g) after additions to ensure all assay components are on the bottom of the wells.
- Cells should be grown to a confluency of 60 to 80% before plating the assay.

**5.2.2 Plate Cells**

1. Harvest cells from culture in Growth Medium and resuspend in Assay Medium to a density of 312,500 cells/ml.
2. Add 32  $\mu$ l per well of Assay Medium to the Cell-free control wells. Add 32  $\mu$ l per well of the cell suspension to Untreated and Compound-treated wells.

**5.2.3 Prepare Stock Solutions**

1. Prepare Assay Medium with 4% DMSO.
2. Prepare 10X Raf1 Inhibitor I in Assay Medium.

**5.2.4 Compound Treatment**

1. Add 4  $\mu$ l Assay Medium with 4% DMSO to all the wells
2. Add 4  $\mu$ l 10X Raf1 Inhibitor I in Assay Medium to the treated wells, and 4  $\mu$ l Assay Medium to the untreated and cell-free control wells
3. Incubate the assay plate in a humidified 37°C/5% CO<sub>2</sub> incubator for 17 hours.

**5.2.5 Substrate Loading and Incubation**

This protocol is designed for loading cells with LiveBLAzer™-FRET B/G Substrate (CCF4-AM) or CCF2-AM. If alternative substrates are used, please follow the loading protocol provided with the substrate.

Preparation of 6X LiveBLAzer™-FRET B/G Substrate (CCF4-AM) or CCF2-AM Mixture and cell loading should be done in the absence of direct strong lighting. Turn off the light in the hood.

1. Prepare Solution A: 1 mM LiveBLAzer™-FRET B/G Substrate (CCF4-AM, MW = 1096) stock solution in dry DMSO. Store the aliquots of the stock solution at -20°C until use.
2. Prepare 6X LiveBLAzer™-FRET B/G (CCF4-AM) Substrate Mixture:
  - 2.1 Add 6  $\mu$ l of Solution A to 60  $\mu$ l of Solution B and vortex.
  - 2.2 Add 874  $\mu$ l Solution C and 60  $\mu$ l of Solution D to the combined solutions from above step with vortexing.
3. Remove assay plate from the humidified 37°C/5% CO<sub>2</sub> incubator.
4. Add 8  $\mu$ l of 6X Substrate Mixture from **Step 2** to each well.
5. Cover the plate to protect it from light and evaporation.
6. Incubate at room temperature for 2.5 hours.

**5.3 Detection**

All measurements using LiveBLAzer™-FRET B/G Substrate are to be made at room temperature from the bottom of the wells. Before reading the plate, remove dust from the bottom with compressed air.

**5.3.1 Instrumentation, Filters, and Plates**

- Fluorescence plate reader with bottom reading capabilities.
- Recommended filters for fluorescence plate reader:
 

Excitation filter:	409/20 nm
Emission filter:	460/40 nm
Emission filter:	530/30 nm

**5.3.2 Reading an Assay Plate**

1. Set the fluorescence plate reader to bottom-read mode.
2. Allow the lamp in the fluorescence plate reader to warm up for at least 10 minutes before making measurements.
3. Use the following filter selections:

	Scan 1	Scan 2
<b>Purpose:</b>	Measure fluorescence in the blue channel	Measure fluorescence in the green channel
<b>Excitation filter:</b>	409/20 nm	409/20 nm
<b>Emission filter:</b>	460/40 nm	530/30 nm

## 6.0 DATA ANALYSIS

### 6.1 Background Subtraction

We recommend that you subtract the background for both emission channels (460 nm and 530 nm).

1. Use the assay plate layout to identify the location of the Cell-free wells. These control wells are used for background subtraction.
2. Determine the average emission from the Cell-free wells at both 460 nm (Average Blue Background) and 530 nm (Average Green Background).
3. Subtract the Average Blue Background (data collected at 460 nm) from all of the blue emission data.
4. Subtract the Average Green background (data collected at 530 nm) from all of the green emission data.
5. Calculate the Blue/Green Emission Ratio for each well, by dividing the background-subtracted blue emission values by the background-subtracted green emission values.

### 6.2 Visual Observation of Intracellular Beta-lactamase Activity Using LiveBLAzer™-FRET B/G Substrate (CCF4-AM)

**Note:** Microscopic visualization of cells will cause photobleaching. Always read the assay plate in the fluorescence plate reader before performing microscopic visualization.

An inverted microscope equipped for epifluorescence and either a xenon or mercury excitation lamp is typically required to view the LiveBLAzer™-FRET B/G Substrate (CCF4-AM) signal in cells. To visually inspect the cells, you will need a long-pass filter passing blue and green fluorescence light so that your eye can visually identify whether the cells are fluorescing green or blue.


Recommended filter sets for observing beta-lactamase activity are described below and are available from Chroma Technologies (800-824-7662, [www.chroma.com](http://www.chroma.com)).

#### Chroma Set # 41031

Excitation filter:	HQ405/20x (405 ± 10 nm)
Dichroic mirror:	425 DCXR
Emission filter:	HQ435LP (435 long-pass)

Filter sizes vary for specific microscopes and need to be specified when the filters are ordered. For epifluorescence microscopes, a long-pass dichroic mirror is needed to separate excitation and emission light and should be matched to the excitation filter (to maximally block the excitation light around 405 nm, yet allow good transmission of the emitted light).

### 6.3 Representative Data



**Figure 1** Dose response of AP1-*bla* A375 cells to Raf1 Inhibitor I. AP1-*bla* A375 cells (10,000 cells/well) were plated in a 384-well format and were treated with the indicated concentrations of Raf1 Inhibitor I (EMD #553008) in the presence of 0.5% DMSO for 17 hours. Cells were then loaded with LiveBLAzer™-FRET B/G Substrate for 2.5 hours. Fluorescence emission values at 460 nm and 530 nm were obtained using a standard fluorescence plate reader and the 460 nm/530 nm ratio plotted for the indicated concentrations of Raf1 Inhibitor I (n=16 for each data point).

## 7.0 DETAILED CELL HANDLING PROCEDURES

### 7.1 Thawing Method

1. Place 14 ml of Thaw Medium (without Blasticidin) into a T75 flask.
2. Place the flask in a humidified 37°C/5% CO<sub>2</sub> incubator for 15 minutes to allow medium to equilibrate to the proper pH and temperature.
3. Remove the vial of cells to be thawed from liquid nitrogen and thaw rapidly by placing at 37°C in a water bath with gentle agitation for 1–2 minutes. Do not submerge vial in water.
4. Decontaminate the vial by wiping with 70% ethanol before opening in a Class II biological safety cabinet.
5. Transfer the vial contents to a sterile 15-ml conical tube.
6. Add 10 ml of Thaw Medium (without blasticidin) dropwise into the cell suspension.
7. Centrifuge cells at 200 × g for 5 minutes.
8. Aspirate supernatant and resuspend the cell pellet in 1 ml of fresh Thaw Medium (without Blasticidin).
9. Transfer contents to the T75 tissue culture flask containing pre-equilibrated Thaw Medium (without Blasticidin) and place flask in a humidified 37°C/5% CO<sub>2</sub> incubator.
10. **At first passage**, switch to Growth Medium with Blasticidin.

### 7.2 Propagation Method

1. Cells should be passaged or fed at least three times a week. Cells should be maintained between 10% and 90% confluency. Do not allow cells to reach confluence.
2. To passage cells, aspirate medium, rinse once with PBS, add Trypsin/EDTA (3 ml for a T75 flask and 5 ml for a T175 flask and 8 ml for a T225 flask) and swirl to coat the cells evenly. Cells usually detach after ~2 minutes exposure to Trypsin/EDTA. Add an equal volume of Growth Medium to inactivate Trypsin. Verify under a microscope that cells have detached and clumps have completely dispersed.
3. Spin down cells and resuspend in Growth Medium

### 7.3 Freezing Method

1. Harvest and count the cells, then spin cells down and resuspend in 4°C Cell Culture Freezing Medium at a density of 2 × 10<sup>6</sup> cells/ml.
2. Dispense 1.0-ml aliquots into cryogenic vials.
3. Place in an insulated container for slow cooling and store overnight at –80°C.
4. Transfer to liquid nitrogen the next day for storage.

## 8.0 REFERENCES

1. Zlokarnik, G., *et al*, **Quantitation of Transcription and Clonal Selection of Single Living Cells with Beta-Lactamase as Reporter**, (1998) *Science*; **279**: p84-88.
2. Kunapuli P., Ransom R., Murphy K., Pettibone D., Kerby J., Grimwood S., Zuck P., Hodder P., Lacson R., Hoffman I., Inglese J., Strulovici B., **Development of an Intact Cell Reporter Gene Beta-lactamase Assay for G Protein-coupled Receptors**, (2003) *Analytical Biochem.*; **314**: p16-29.
3. Xing, H., Pollok, B., *et al*, **A Fluorescent Reporter Assay For The Detection of Ligands Acting Through G1 Protein-coupled Receptors**, (2000) *J. Receptor & Signal Transduction Research*; **20**: p189-210.
4. Qureshi, S., *et al*, **A One-Arm Homologous Recombination Approach for Developing Nuclear Receptor Assays in Somatic Cells**, (2003) *Assay and Drug Dev. Tech*; **1**: p755-766.
5. Peekhaus, N. *et al*, **A Beta-Lactamase-Dependent Gal4-Estrogen Receptor Transactivation Assay for the Ultra-High Throughput Screening of Estrogen Receptor Agonists in a 3,456-Well Format**, (2003) *Assay and Drug Dev Tech*; **1**: p789-800.
6. Chin, J., *et al*, **Miniaturization of Cell-Based Beta-Lactamase-Dependent FRET Assays to Ultra-High Throughput Formats to Identify Agonists of Human Liver X Receptors**, (2003) *Assay and Drug Dev. Tech.*; **1**: p777-787.
7. Whitney M., Rockenstein E., Cantin G., Knapp T., Zlokarnik G., Sanders P., Durick K., Craig F.F., Negulescu P.A., **A Genome-wide Functional Assay of Signal Transduction in Living Mammalian Cells**, (1998) *Nat. Biotechnol.*; **16**: p1329-1333.

## 9.0 PURCHASER NOTIFICATION

### Limited Use Label License No. 5: Invitrogen Technology

The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes. The buyer may transfer information or materials made through the use of this product to a scientific collaborator, provided that such transfer is not for any Commercial Purpose, and that such collaborator agrees in writing (a) not to transfer such materials to any third party, and (b) to use such transferred materials and/or information solely for research and not for Commercial Purposes. Commercial Purposes means any activity by a party for consideration and may include, but is not limited to: (1) use of the product or its components in manufacturing; (2) use of the product or its components to provide a service, information, or data; (3) use of the product or its components for therapeutic, diagnostic or prophylactic purposes; or (4) resale of the product or its components, whether or not such product or its components are resold for use in research. Invitrogen Corporation will not assert a claim against the buyer of infringement of patents owned or controlled by Invitrogen Corporation which cover this product based upon the manufacture, use or sale of a therapeutic, clinical diagnostic, vaccine or prophylactic product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. If the purchaser is not willing to accept the limitations of this limited use statement, Invitrogen is willing to accept return of the product with a full refund. For information on purchasing a license to this product for purposes other than research, contact Licensing Department, Invitrogen Corporation, 1600 Faraday Avenue, Carlsbad, California 92008. Phone (760) 603-7200. Fax (760) 602-6500. Email: [outlicensing@invitrogen.com](mailto:outlicensing@invitrogen.com).

### Limited Use Label License No. 150: GeneBLAzer® Technology

This product and/or its use is the subject of one or more of U.S. Patent Nos. 5,741,657, 5,955,604, 6,291,162, and 6,472,205 and foreign equivalents licensed to Invitrogen Corporation. The right to use this product for internal research specifically excludes the right to use this product to identify, discover, and profile compounds that act as a flavor, fragrance or taste-enhancers and modify a target identified in taste, olfaction, or pheromone detection, which compound does not require FDA approval of an NDA for claims of safety and efficacy. The right to use methods claimed in the foregoing patents with this product for research purposes can only be acquired by the use of GeneBLAzer® substrates purchased from Invitrogen Corporation or its authorized distributors.

### Limited Use Label License No. 51: Blasticidin and the Blasticidin Selection Marker

Blasticidin and the blasticidin resistance gene (*bsd*) are the subject of U.S. Patent No. 5,527,701 sold under patent license for research purposes only. For information on purchasing a license to this product for purposes other than research, contact Licensing Department, Invitrogen Corporation, 1600 Faraday Avenue, Carlsbad, California 92008. Phone (760) 603-7200. Fax (760) 602-6500.

### Use of Genetically Modified Organisms (GMO)

**Information for European Customers** The CellSensor® AP1-*bla* A375 cell line(s) are genetically modified with the plasmid pLenti-*bsd* / AP1-*bla*. As a condition of sale, use of this product must be in accordance with all applicable local legislation and guidelines including EC Directive 90/219/EEC on the contained use of genetically modified organisms.

© 2007, Invitrogen Corporation. All rights reserved. Reproduction forbidden without permission.