



Miltenyi Biotec

# StemMACS™ PD0325901

2 mg  
5×2 mg

130-103-923  
130-104-170

## Contents

1. Description
  - 1.1 Background information
2. Protocol
  - 2.1 Preparation of stock solution
  - 2.2 Use in cell culture

## 1. Description

**Components** StemMACS™ PD0325901. A selective small molecule MEK inhibitor.

**Size** 2 mg; 5×2 mg

**Product format** Off-white solid


**Molecular weight** 482.19

**CAS number** 391210-10-9

**Systematic name** N-[(2R)-2,3-dihydroxypropoxy]-3,4-difluoro-2-[(2-fluoro-4-iodophenyl)amino]-benzamide

**Molecular formula** C<sub>16</sub>H<sub>14</sub>F<sub>3</sub>IN<sub>2</sub>O<sub>4</sub>

### Structure



**Purity** >98% by LC/MS

**Solubility** Soluble in DMSO (up to 25 mM)

**Storage** Store powder at -20 °C. After reconstitution, store aliquots at -20 °C. Protect from light.

**Quality control** Purity and identity of StemMACS PD0325901 were determined by LC/MS. Cell culture compatibility was tested on pluripotent stem cell cultures.

### 1.1 Background information

StemMACS™ PD0325901 is a selective small molecule inhibitor of MEK (mitogen-activated protein kinase, MAPK/ERK kinase), a key component of the Ras/Raf/MEK/ERK signaling pathway. PD0325901 is a derivative of the MEK inhibitor CI-1040 and shows improved cell permeability and higher potency towards ERK1/2 (IC<sub>50</sub>=0.33 nM). In pluripotent stem cell research, inhibition of the MAPK signaling pathway with PD0325901 has been used to enhance reprogramming and sustain pluripotent stem cell self-renewal.

## 2. Protocol

### 2.1 Preparation of stock solution

Effective concentrations of StemMACS PD0325901 for cell culture applications range from 0.1 μM to 2 μM. A 10 mM stock solution in DMSO will be appropriate for most applications and can be prepared as follows:

1. Reconstitute the entire vial contents by adding 414.8 μL of pure DMSO. Warm to 37 °C for 3–5 minutes to facilitate solubilization.

▲ **Note:** The vial may have turned upside down during transportation. Gently tap prior to reconstitution to collect all powder at the bottom of the vial.

2. Prepare appropriate aliquots and store at -20 °C. Avoid repeated freeze-thaw cycles.

▲ **Note:** The DMSO concentration in culture should not exceed 0.5%. Stock solutions of alternate concentration can be prepared using the following table. Add the solvent directly to the vial, it will hold up to 4 mL.

Desired stock	1 mM	2 mM	5 mM	10 mM	20 mM
Volume of DMSO to add	Dilute 1:10 from a 10 mM stock	2074 μL	829.6 μL	414.8 μL	207.4 μL

### 2.2 Use in cell culture

1. Thaw aliquots at 37 °C as needed.
2. To avoid precipitation, prewarm the cell culture media prior to adding the reconstituted compound.
3. Mix and filter the supplemented media through a 0.2 μM low-protein binding filter.

All protocols and data sheets are available at [www.miltenyibiotec.com](http://www.miltenyibiotec.com).

### Warranty

The products sold hereunder are warranted only to be free from defects in workmanship and material at the time of delivery to the customer. Miltenyi Biotec GmbH makes no warranty or representation, either expressed or implied, with respect to the fitness of a product for a particular purpose. There are no warranties, expressed or implied, which extend beyond the technical specifications of the products. Miltenyi Biotec GmbH's liability is limited to either replacement of the products or refund of the purchase price. Miltenyi Biotec GmbH is not liable for any property damage, personal injury or economic loss caused by the product.

MACS is a registered trademark and StemMACS is a trademark of Miltenyi Biotec GmbH.

Unless otherwise specifically indicated, Miltenyi Biotec products and services are for research use only and not for diagnostic or therapeutic use.

Copyright © 2014 Miltenyi Biotec GmbH. All rights reserved.