

## Contents


1. Description
  - 1.1 Principle of the staining procedure
  - 1.2 Background information
  - 1.3 Applications
  - 1.4 Reagent requirements
2. General protocol for fluorescent staining
3. Example of a fluorescent staining with Annexin V-FITC Kit
4. References

## 1. Description

<b>Components</b>	<b>1 mL Annexin V-FITC:</b> Annexin V conjugated to FITC (fluorescein-isothiocyanate) <b>25 mL 20× Binding Buffer Stock Solution</b> <b>0.5 mL Propidium Iodide (100 µg/mL)</b>
<b>Capacity</b>	For 10 <sup>8</sup> total cells, up to 100 stainings.
<b>Product format</b>	Annexin V-FITC is supplied in buffer containing stabilizer and 0.05% sodium azide.
<b>Storage</b>	Store protected from light at 2–8 °C. Do not freeze. The expiration date is indicated on the vial label.

### 1.1 Principle of the staining procedure

The Annexin V-FITC Kit has been developed for detection and discrimination of apoptotic, necrotic, and dead cells. Apoptotic cells are stained positively for Annexin V-FITC that binds to phosphatidylserine (PS), but are negative for staining with propidium iodide (PI). Dead cells are stained positive for Annexin V-FITC and PI, whereas viable cells are negative for both Annexin V-FITC and PI.



## 1.2 Background information

In most normal, viable eukaryotic cells, the negatively charged phospholipid phosphatidylserine (PS) is located in the cytosolic leaflet of the plasma membrane lipid bilayer.<sup>1</sup> PS redistribution from the inner to the outer leaflet is an early and widespread event during apoptosis.<sup>1,2</sup> However, in necrosis, PS becomes accessible due to the disruption of membrane integrity.<sup>2</sup> Apart from necrosis and apoptosis, PS also becomes accessible in activated platelets<sup>3</sup>, in certain cell anomalies like sickle cell anemia<sup>4</sup>, in erythrocyte senescence<sup>5</sup>, upon degranulation of mast cells<sup>6</sup> and in certain stages of B cell differentiation<sup>7</sup>. PS exposure also serves as a trigger for the recognition and removal of apoptotic cells by macrophages.<sup>8,9</sup> Annexin V is a 35 kDa phospholipid-binding protein and a major cell membrane component of macrophages and other phagocytic cell types. Annexin V has a high affinity to PS in the presence of physiological concentrations of calcium (Ca<sup>2+</sup>).<sup>10</sup>

## 1.3 Applications

- Studies on cell death (apoptosis and/or necrosis).
- Evaluation of separations with the Annexin V MicroBead Kit (# 130-090-201).

## 1.4 Reagent and instrument requirements

- Buffer (1× Binding Buffer)  
 Prepare 1× Binding Buffer from 20× Binding Buffer Stock Solution by diluting 250 µL of 20× Binding Buffer Stock Solution with 4.75 mL of sterile, distilled water. This volume is sufficient for 10<sup>6</sup> total cells. Alternatively, prepare 1× Binding Buffer Stock Solution by diluting 25 mL of 20× Binding Buffer with 475 mL of sterile, distilled water.  
 Store at 4 °C.

▲ **Note:** Handle under sterile conditions.

## 2. General protocol for fluorescent staining

▲ Volumes for fluorescent labeling given below are for up to 10<sup>6</sup> total cells. When working with fewer than 10<sup>6</sup> cells, use the same volumes as indicated. When working with higher cell numbers, scale up all reagent volumes and total volumes, accordingly (e.g. for 2×10<sup>6</sup> total cells, use twice the volume of all indicated reagent volumes and total volumes).


▲ When working with cell samples containing platelets, for example, blood samples, wash samples carefully at 200×g in order to remove platelets. Use buffer containing the ion chelator EDTA for these washing steps. Activated platelets expose PS and therefore bind Annexin V.<sup>5</sup>

1. Determine cell number.
2. Wash 10<sup>6</sup> cells in 1 mL of 1× Binding Buffer and centrifuge at 300×g for 10 minutes. Aspirate supernatant completely.
3. (Optional) Repeat washing step.

4. Resuspend cell pellet in 100  $\mu$ L of 1 $\times$  Binding Buffer per  $10^6$  cells.
5. Add 10  $\mu$ L of Annexin V-FITC per  $10^6$  cells.
6. Mix well and incubate for 15 minutes in the dark at room temperature.
7. Wash cells by adding 1 mL of 1 $\times$  Binding Buffer per  $10^6$  cells and centrifuge at 300 $\times$ g for 10 minutes. Aspirate supernatant completely.
8. (Optional) Repeat washing step.
9. Resuspend cell pellet in 500  $\mu$ L of 1 $\times$  Binding Buffer per  $10^6$  cells.
10. Add 5  $\mu$ L of PI solution immediately prior to analysis by flow cytometry or fluorescence microscopy.

### 3. Example of a fluorescent staining with Annexin V-FITC Kit

Jurkat cells cultured with staurosporine (50 nM) for 15 h, were stained with Annexin V-FITC and PI and analyzed by flow cytometry.



### 4. References

1. Koopman, G. *et al.* (1994) Annexin V for Flow Cytometric Detection of Phosphatidylserine expression on B Cells Undergoing Apoptosis. *Blood* 84: 1415–1420.
2. Martin, S. J. *et al.* (1995) Early Redistribution of Plasma Membrane Phosphatidylserine Is a General Feature of Apoptosis Regardless of the Initiating Stimulus: Inhibition by Overexpression of Bcl-2 and Abl. *J. Exp. Med.* 182: 1545–1556.
3. Thiagarajan, P. and Tait, J. F. (1990) Binding of annexin V/placental anticoagulant protein I to platelets. Evidence for phosphatidylserine exposure in the procoagulant response of activated platelets. *J Biol Chem* 265: 17420–17423.
4. Kuypers, F. A. *et al.* (1996) Detection of Altered Membrane Phospholipid Asymmetry in Subpopulations of Human Red Blood Cells Using Fluorescently Labeled Annexin V. *Blood* 87: 1179–1187. [181]
5. Schroit, A. J. and Zwaal, R. F. A. (1991) Transbilayer movement of phospholipids in red cells and platelet membranes. *Biochim. Biophys. Acta.* 1071: 313–329.
6. Demo, S. D. *et al.* (1999) Quantitative Measurement of Mast Cell Degranulation Using a Novel Flow Cytometric Annexin-V Binding Assay. *Cytometry* 36: 340–348.
7. Dillon, S. R. *et al.* (2001) Annexin V Binds to Positively Selected B Cells. *J. Immunol.* 166: 58–71.
8. Fadok, V. A. *et al.* (1992) Exposure of Phosphatidylserine on the Surface of Apoptotic Lymphocytes Triggers Specific Recognition and Removal by Macrophages. *J. Immunol.* 148: 2207–2216.
9. Fadok, V. A. *et al.* (2000) A receptor for phosphatidylserine-specific clearance of apoptotic cells. *Nature* 405: 85–90.
10. Moss, S. E. *et al.* (1991) Diversity in the Annexin Family. In *Novel Calcium Binding Proteins*, Springer Verlag, 535–566.

All protocols and data sheets are available at [www.miltenyibiotec.com](http://www.miltenyibiotec.com).

### Warnings

Reagents contain sodium azide. Under acidic conditions sodium azide yields hydrazoic acid, which is extremely toxic. Azide compounds should be diluted with running water before discarding. These precautions are recommended to avoid deposits in plumbing where explosive conditions may develop.

### Warranty

The products sold hereunder are warranted only to be free from defects in workmanship and material at the time of delivery to the customer. Miltenyi Biotec GmbH makes no warranty or representation, either expressed or implied, with respect to the fitness of a product for a particular purpose. There are no warranties, expressed or implied, which extend beyond the technical specifications of the products. Miltenyi Biotec GmbH's liability is limited to either replacement of the products or refund of the purchase price. Miltenyi Biotec GmbH is not liable for any property damage, personal injury or economic loss caused by the product.

MACS is a registered trademark of Miltenyi Biotec GmbH.

Copyright © 2009 Miltenyi Biotec GmbH. All rights reserved.