



**Rabbit (polyclonal)
Anti-GATA-4 [pS¹⁰⁵]
Phosphospecific Antibody, Unconjugated**

PRODUCT ANALYSIS SHEET

Catalog Number/Size:	44-948 (10 mini-blot size)
Lot Number:	See product label
Volume:	100 µL
Concentration:	See product label
Form of Antibody:	Rabbit polyclonal immunoglobulin in Dulbecco's phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.3 (+/- 0.1), with 1.0 mg/mL BSA (IgG, protease free) as a carrier.
Preservative:	0.05% sodium azide (Caution: sodium azide is a poisonous and hazardous substance. Handle with care and dispose of properly.)
Purification:	Purified from rabbit serum by sequential epitope-specific chromatography. The antibody has been negatively preadsorbed using a non-phosphopeptide corresponding to the site of phosphorylation to remove antibody that is reactive with non-phosphorylated GATA-4. The final product is generated by affinity chromatography using a GATA-4 derived peptide that is phosphorylated at serine 105.
Immunogen:	The antiserum was produced against a chemically synthesized phosphopeptide derived from a region of human GATA-4 that contains serine 105. The sequence is conserved in mouse, rat and chicken.
Target Summary:	GATA-4 is a 46 kDa member of the GATA family of zinc finger-containing transcription factors that is involved in the development of cardiac hypertrophy and remodeling, and plays a critical role in regulating basal and agonist or stress-induced gene expression in cardiac and smooth muscle cell types. GATA-4 contains a conserved MAPK phosphorylation site at serine 105 within the transcriptional activation domain. Serine 105 of GATA-4 is phosphorylated in response to agonist stimulation through MEK1-ERK1/2, and weakly through JNK or p38 MAPKs.
Reactivity:	Mouse GATA-4. Human, rat and chicken (100% homologous) GATA-4 have not been tested, but are expected to react.
Applications:	The antibody has been used in Western blotting. Other applications have not been tested at Invitrogen.
Suggested Working Dilutions:	For Western blotting applications, we recommend using the antibody at 0.1-1.0 µg/mL. At 0.35 µg/mL, the dilution provides 100 mL working solution, which at 10 mL/blot allows 10 blots to be performed. The optimal antibody concentration should be determined empirically for each specific application.
Storage:	Store at -80°C. Upon initial thawing, apportion into working aliquots and store at -80°C. Avoid repeated freeze-thaw cycles to prevent denaturing the antibody.
Expiration Date:	Expires one year from date of receipt when stored as instructed.
Positive Controls Used:	Phenylephrine (PE)-stimulated rat cardiomyocytes, transfected with mouse GATA-4.

This product is for research use only. Not for use in diagnostic procedures.

www.invitrogen.com

Invitrogen Corporation • 542 Flynn Rd • Camarillo • CA 93012 • Tel: 800.955.6288 • E-mail: techsupport@invitrogen.com
This antibody is manufactured under a licensed process covered by Patent # 5, 599, 681.

PI44948

(Rev 11/08) DCC-08-1089

Important Licensing Information - These products may be covered by one or more Limited Use Label Licenses (see the Invitrogen Catalog or our website, www.invitrogen.com). By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

Related Products:**Antibodies:**

c-Met [pYpYpY^{1230/1234/1235}], Cat. # 44-888G ERK1&2 [pTpY^{185/187}], Cat. # 44-680G
GSK3-β [pS⁹], Cat. # 44-600G JNK1&2 [pTpY^{183/185}], Cat. # 44-682G
PLCγ-1 [pY⁷⁸³], Cat. # 44-696G p38 [pTpY^{180/182}], Cat. # 44-684G
NFAT-1 [pS⁵⁴] (mouse), Cat. # 44-944G PKCα pan, Cat. # AHO0622


References:

Kitta, K., et al. (2002) Hepatocyte growth factor induces GATA-4 phosphorylation and cell survival in cardiac muscle cells. *J. Biol. Chem.* Dec. 4 [epub ahead of print].

Clement, S.A., et al. (2002) Roles of protein kinase C and alpha-tocopherol in regulation of signal transduction for GATA-4 phosphorylation in HL-1 cardiac muscle cells. *Free Radic. Biol. Med.* 32(4):341-349.

Liang, Q., et al. (2001) The transcription factor GATA-4 is activated by extracellular signal-regulated kinase 1- and 2- mediated phosphorylation of serine 105 in cardiomyocytes. *Mol. Cell. Biol.* 21(21):7460-7469.

Morisco, C., et al. (2001) Glycogen synthase kinase 3beta regulates GATA-4 in cardiac myocytes. *J. Biol. Chem.* 276(30):28586-28597.

**Mutant Analysis and Antibody-Peptide Competition**

Extracts of cardiomyocytes overexpressing mutant GATA-4 S105A (1) or wild-type GATA-4 (2-5) stimulated with phenylephrine (PE) for 24 hours were resolved by SDS-PAGE on a 10% Tris-glycine gel and transferred to PVDF. The membrane was blocked with a 5% BSA-TBST buffer overnight at 4°C, then incubated with 0.35 µg/mL GATA-4 [pS¹⁰⁵] antibody for two hours at room temperature in a 3% BSA-TBST buffer, following prior incubation with: no peptide (1, 2), the non-phosphopeptide corresponding to the phosphopeptide immunogen (3), a generic phosphoserine-containing peptide (4), or the phosphopeptide immunogen (5). After washing, the membrane was incubated with goat F(ab')₂ anti-rabbit IgG alkaline phosphatase (cat.# ALI4405) and signals were detected using the Tropix WesternStar™ method.

The data show that only the phosphopeptide corresponding to GATA-4 [pS¹⁰⁵] blocks the antibody signal, and that the antibody does not recognize the GATA-4 S105A point mutant, demonstrating the specificity of the antibody.

This product is for research use only. Not for use in diagnostic procedures.

www.invitrogen.com

Invitrogen Corporation • 542 Flynn Rd • Camarillo • CA 93012 • Tel: 800.955.6288 • E-mail: techsupport@invitrogen.com

This antibody is manufactured under a licensed process covered by Patent # 5, 599, 681.

PI44948

(Rev 11/08) DCC-08-1089

Important Licensing Information - These products may be covered by one or more Limited Use Label Licenses (see the Invitrogen Catalog or our website, www.invitrogen.com). By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

Western Blotting Procedure

1. Lyse approximately 10^7 cells in 0.5 mL of ice cold Cell Lysis Buffer (formulation provided below). This buffer, a modified RIPA buffer, is suitable for recovery of most proteins, including membrane receptors, cytoskeletal-associated proteins, and soluble proteins. This cell lysis buffer formulation is available as a separate product which requires supplementation with protease inhibitors immediately prior to use (Invitrogen catalog number FNN0011). Other cell lysis buffer formulations, such as Laemmli sample buffer and Triton-X 100 buffer, are also compatible with this procedure. Additional optimization of the cell stimulation protocol and cell lysis procedure may be required for each specific application.
2. Remove the cellular debris by centrifuging the lysates at 14,000 x g for 10 minutes. Alternatively, lysates may be ultracentrifuged at 100,000 x g for 30 minutes for greater clarification.
3. Carefully decant the clarified cell lysates into clean tubes and determine the protein concentration using a suitable method, such as the Bradford assay. Polypropylene tubes are recommended for storing cell lysates.
4. React an aliquot of the lysate with an equal volume of 2x Laemmli Sample Buffer (125 mM Tris, pH 6.8, 10% glycerol, 10% SDS, 0.006% bromophenol blue, and 130 mM dithiothreitol [DTT]) and boil the mixture for 90 seconds at 100°C.
5. Load 10-30 µg of the cell lysate into the wells of an appropriate single percentage or gradient minigel and resolve the proteins by SDS-PAGE.
6. In preparation for the Western transfer, cut a piece of PVDF membrane slightly larger than the gel. Soak the membrane in methanol for 1 minute, then rinse with ddH₂O for 5 minutes.
7. Soak the membrane, 2 pieces of Whatman paper, and Western apparatus sponges in transfer buffer (formulation provided below) for 2 minutes.
8. Assemble the gel and membrane into the sandwich apparatus.
9. Transfer the proteins at 140 mA for 60-90 minutes at room temperature.
10. Following the transfer, rinse the membrane with Tris buffered saline for 2 minutes.
11. Block the membrane with blocking buffer (formulation provided below) overnight at 4°C or for one hour at room temperature.
12. Incubate the blocked blot with primary antibody at a concentration of 0.1-1.0 µg/mL in Tris buffered saline supplemented with 3% Ig-free BSA and 0.1% Tween 20 overnight at 4°C or for 2 hours at room temperature.
13. Wash the blot with several changes of Tris buffered saline supplemented with 0.1% Tween 20.
14. Detect the antibody band using an appropriate secondary antibody, such as goat F(ab')₂ anti-rabbit IgG alkaline phosphatase conjugate (catalog number ALI4405) or goat F(ab')₂ anti-rabbit IgG horseradish peroxidase conjugate (catalog number ALI4404) in conjunction with your chemiluminescence reagents and instrumentation.

Cell Lysis Buffer

Formulation:

10 mM Tris, pH 7.4
100 mM NaCl
1 mM EDTA
1 mM EGTA
1 mM NaF
20 mM Na₄P₂O₇
2 mM Na₃VO₄
0.1% SDS
0.5% sodium deoxycholate
1% Triton-X 100
10% glycerol
1 mM PMSF (made from a
0.3 M stock in DMSO)
or 1 mM AEBSF (water
soluble version of PMSF)
60 µg/mL aprotinin
10 µg/mL leupeptin
1 µg/mL pepstatin
(alternatively, protease inhibitor
cocktail such as Sigma catalog
number P2714 may be used)

Transfer Buffer

Formulation:

2.4 gm Tris base
14.2 gm glycine
200 mL methanol
Q.S. to 1 liter, then add
1 mL 10% SDS.
Cool to 4°C prior to use.

Tris Buffered Saline

Formulation:

20 mM Tris-HCl, pH 7.4
0.9% NaCl

Blocking Buffer

Formulation:

100 mL Tris buffered saline
5 gm Ig-free BSA
0.1 mL Tween 20

This product is for research use only. Not for use in diagnostic procedures.

www.invitrogen.com

Invitrogen Corporation • 542 Flynn Rd • Camarillo • CA 93012 • Tel: 800.955.6288 • E-mail: techsupport@invitrogen.com

This antibody is manufactured under a licensed process covered by Patent # 5, 599, 681.

PI44948

(Rev 11/08) DCC-08-1089

Important Licensing Information - These products may be covered by one or more Limited Use Label Licenses (see the Invitrogen Catalog or our website, www.invitrogen.com). By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

Peptide Competition Experiment

To demonstrate the specificity of a Phosphorylation Site Specific Antibody, we recommend the following peptide competition experiment which uses our control peptides. These control peptides have the sequences of the phosphopeptide immunogen used to raise the antibody and the corresponding non-phosphorylated peptide. In the competition experiment, 200-500 fold molar excess of the phosphorylated and non-phosphorylated peptides are pre-incubated with aliquots of the antibody prior to use in immunoassay procedures.

A sample calculation for the determination of the 200 fold molar excess of peptide to antibody is presented below. The following assumptions have been made:

- The molecular mass of an IgG molecule is 150,000 daltons.
- Each mole of antibody binds two moles of peptide.
- The Phosphorylation Site Specific Antibody is used at a concentration of 0.5 µg/mL.

The optimal antibody concentration for use in peptide competition experiments is below saturating as determined by previous experiments in your system. If an optimal concentration has not been determined, it is suggested that the concentration provided on the antibody Product Analysis Sheet be used. A final antibody concentration of 0.5 µg/mL is satisfactory for most applications.

The molarity of the 0.5 µg/mL antibody solution is:

$$(0.5 \mu\text{g/mL})(1000 \text{ mL/L})/(150,000 \mu\text{g}/\mu\text{mole}) = 0.00333 \mu\text{M}.$$

Because each mole of antibody binds two moles of peptide, 0.5 µg/mL antibody can bind 0.00667 µM of peptide.

A 200 fold molar excess of peptide is $(200)(0.00667 \mu\text{M}) = 1.334 \mu\text{M}$.

The following procedure describes peptide competition experiments using antibody at a concentration of 0.5 µg/mL and a 200 fold molar excess of peptides based on the calculation above, in a total volume of 2 mL.

Procedure:

1. Prepare three identical test samples, such as identical nitrocellulose or PVDF strips with transferred protein. The test samples should be blocked with BSA or non-fat dried milk in a buffer compatible with an antibody based detection method, such as Tris buffered saline or phosphate buffered saline.
2. Slowly thaw the Phosphorylation Site Specific Antibody on ice.
3. Prepare 3 mL of a 2x (1 µg/mL) antibody stock solution in a buffer appropriate for the application. Suggested buffer formulations are TBS or PBS supplemented with blocking protein such as BSA or non-fat dried milk.
4. Apportion the unused Phosphorylation Site Specific Antibody into working aliquots and store at -80°C for future use.
5. The lyophilized control peptides should be warmed to room temperature, ideally under desiccation.
6. Reconstitute each of the control peptides to a concentration of 100 µM using nanopure water at room temperature. As indicated on the peptide labels, each vial contains 0.1 mg. For a peptide with a molecular mass of 1500, reconstitution with 0.67 mL water yields a solution with a concentration of 100 µM.
7. Allow the peptides to dissolve at room temperature, then gently triturate several times using a pipette. Avoid introducing air bubbles.
8. Label 3 test tubes as follows:
 - tube 1: water only no peptide control
 - tube 2: phosphopeptide
 - tube 3: non-phosphopeptide
9. Prepare 2x peptide stock solutions (2.66 µM) or water control by pipetting the following:
 - tube 1: water control stock: 0.973 mL buffer (TBS or PBS plus blocking protein) plus 27 µL water.
 - tube 2: phosphopeptide 2x stock: 0.973 mL buffer (TBS or PBS plus blocking protein) plus 27 µL reconstituted (100 µM) phosphopeptide.
 - tube 3: non-phosphopeptide 2x stock: 0.973 mL buffer (TBS or PBS plus blocking protein) plus 27 µL reconstituted (100 µM) non-phosphopeptide.
10. Apportion unused reconstituted peptide solutions into working aliquots and store at -20°C for future use.
11. Pipette 1 mL of the 2x antibody stock into each of the tubes marked 1, 2, and 3. The tubes should be incubated for 30 minutes at room temperature with gentle rocking.
12. The pre-incubated antibody in each of the three tubes is then ready for use. Pipette the contents of each tube onto the three identical test samples.

For Western blotting strips:

 - ♦ Incubate these strips for 2 hours at room temperature, followed by several washes to remove unbound antibody.
 - ♦ Transfer each strip to a new solution containing a labeled secondary antibody (example goat anti-rabbit IgG-alkaline phosphatase conjugate).
 - ♦ Remove unbound secondary antibody by thorough washing and develop bands.

The signals obtained with antibody incubated with “(1) water only no peptide control”, which represents the maximum signal, and the signals obtained with “(2) phosphopeptide and “(3) non-phosphopeptide” are readily compared under these conditions.

This product is for research use only. Not for use in diagnostic procedures.

www.invitrogen.com

Invitrogen Corporation • 542 Flynn Rd • Camarillo • CA 93012 • Tel: 800.955.6288 • E-mail: techsupport@invitrogen.com

This antibody is manufactured under a licensed process covered by Patent # 5, 599, 681.

PI44948

(Rev 11/08) DCC-08-1089

Important Licensing Information - These products may be covered by one or more Limited Use Label Licenses (see the Invitrogen Catalog or our website, www.invitrogen.com). By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.