



04-6111 1 ml
FITC-Rat Anti-
Mouse IgG1
Lot No. See product label

FITC-Monoclonal Rat Anti-Mouse IgG1
For Research Use Only

INVITROGEN MONOCLONAL ANTIBODY REAGENT

FORM:

Liquid. The product is in 10 mM phosphate buffered saline, pH 7.4, containing 1% bovine serum albumin, 40% glycerol, and 0.1% sodium azide as a preservative.

PURIFIED MONOCLONAL RAT ANTI-MOUSE IgG1 SPECIFIC ANTIBODY is conjugated with fluorescein isothiocyanate (FITC) under optimum conditions and is free of unconjugated FITC.

SUGGESTED DILUTION:

1:20 with PBS containing 10% normal rat serum. Working dilution for specific application should be determined by the investigator to obtain the best conditions.

APPLICATIONS:

Fluorescent-antibody conjugates can be used to localize or to label antigens in cells, tissue sections or other fluorescent immunoassays.

STORAGE:

Store at 4°C.

www.invitrogen.com

Invitrogen Corporation • 542 Flynn Rd • Camarillo • CA 93012 • Tel: 800.955.6288

E-mail: techsupport@invitrogen.com

PI046111

(Rev 10/08) DCC-08-1089

Important Licensing Information - These products may be covered by one or more Limited Use Label Licenses (see the Invitrogen Catalog or our website, www.invitrogen.com). By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.



04-6111 1 ml
FITC-Rat Anti-
Mouse IgG1
Lot No. See product label

FITC-Monoclonal Rat Anti-Mouse IgG1
For Research Use Only

INVITROGEN MONOCLONAL ANTIBODY REAGENT

FORM:

Liquid. The product is in 10 mM phosphate buffered saline, pH 7.4, containing 1% bovine serum albumin, 40% glycerol, and 0.1% sodium azide as a preservative.

PURIFIED MONOCLONAL RAT ANTI-MOUSE IgG1 SPECIFIC ANTIBODY is conjugated with fluorescein isothiocyanate (FITC) under optimum conditions and is free of unconjugated FITC.

SUGGESTED DILUTION:

1:20 with PBS containing 10% normal rat serum. Working dilution for specific application should be determined by the investigator to obtain the best conditions.

APPLICATIONS:

Fluorescent-antibody conjugates can be used to localize or to label antigens in cells, tissue sections or other fluorescent immunoassays.

STORAGE:

Store at 4°C.

www.invitrogen.com

Invitrogen Corporation • 542 Flynn Rd • Camarillo • CA 93012 • Tel: 800.955.6288

E-mail: techsupport@invitrogen.com

PI046111

(Rev 10/08) DCC-08-1089

Important Licensing Information - These products may be covered by one or more Limited Use Label Licenses (see the Invitrogen Catalog or our website, www.invitrogen.com). By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.