



Qty: 100 µg/200 µl

Mouse anti-Desmocollin-2/3

Catalog No. 32-6200

Lot No.

Mouse anti-Desmocollin-2/3

FORM

This monoclonal antibody is supplied as a 200 µl aliquot at a concentration of 0.5 mg/ml in PBS, pH 7.4, containing 0.1% sodium azide. This antibody is highly purified from mouse ascites by protein A affinity chromatography.

CLONE: 7G6

ISOTYPE: Mouse IgG₁

IMMUNOGEN

Human desmocollin-2 extracellular domain.

SPECIFICITY

This antibody reacts with the ~120 kDa desmocollin-2 and ~105 kDa desmocollin-3 protein, with particularly strong reaction to desmocollin-2. Cross reactivity with other desmocollin family members has not been observed.

REACTIVITY

This antibody is confirmed reactive with human. Reactivity with other species has not been tested. Positive control: A431 cells.

Sample	Immuno-precipitation (native)	Western Blotting (ECL)	Immuno-fluorescence
Human	+++	+++	+++
Immunogen			

(Excellent +++, Good ++, Poor +, No reactivity 0, Not applicable NA)

USAGE

Working concentrations for specific applications should be determined by the investigator. Appropriate concentrations will be affected by several factors, including secondary antibody affinity, antigen concentration, sensitivity of detection method, temperature and length of incubations, etc. The suitability of this antibody for applications other than those listed below has not been determined. The following concentration ranges are recommended starting points for this product.

Immunoprecipitation: 5-10 µg

Western Blotting: 1-3 µg/ml

Immunofluorescence: 20-50 µg/ml

STORAGE

Store at 2-8°C for up to one month. Store at -20°C for long term storage. Avoid repeated freezing and thawing.

(cont'd)

www.invitrogen.com

Invitrogen Corporation • 542 Flynn Rd • Camarillo • CA 93012 • Tel: 800.955.6288 • E-mail: techsupport@invitrogen.com

PI326200

(Rev 10/08) DCC-08-1089

Important Licensing Information - These products may be covered by one or more Limited Use Label Licenses (see the Invitrogen Catalog or our website, www.invitrogen.com). By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BACKGROUND

Desmosomes are intercellular adhering junctions that represent cell surface attachment sites for intermediate filament. The desmosome is subdivided into two regions. The plaque region lies adjacent to the plasma, and is believed to contain molecules that attach the intermediate filament cytoskeleton to the desmosome. The core region is composed of transmembrane glycoproteins that are thought to mediate cell-cell adhesion.

Desmogleins and desmocollins are the main desmosomal transmembrane proteins. These desmosomal glycoproteins belong to the members of the cadherin family of adhesion molecules. Three different isoforms of both desmogleins and desmocollins have been identified, named as desmoglein1-3 and desmocollin1-3⁽¹⁾. Desmosomal cadherins showed differentiation-specific expression in the human epidermis, although the functional significance of this differential expression is not fully understood. Desmocollin-1 can be found in the upper layers⁽²⁾. The expression of desmocollin-2 varies in the basal and suprabasal layers⁽²⁾. And desmocollin-3 is expressed more evenly throughout the suprabasal layers⁽²⁾.

Domain specific desmosomal and adherens junction-associated antibodies have been widely used in recent studies. In one study, domain specific antibodies were used to determine the cellular localization of the skin of patients with Hailey-Hailey disease and Darier's disease⁽³⁾. Desmosomal cadherin antibodies are also useful for targeting specific disease⁽⁴⁾ and in the study of interaction between cell adhesion proteins⁽⁵⁾.

REFERENCES

1. Buxton, R.S., et al. Nomenclature of the desmosomal cadherins. *J Cell Biol.* 121:481-483 (1993).
2. Kurzen, H., et al. Compositionally different desmosomes in the various compartments of the human hair follicle. *Differentiation* 63(5):295-304 (1998).
3. Hakuno, M., et al. Dissociation of intra- and extracellular domains of desmosomal cadherins and E-cadherin in Hailey-Hailey disease and Darier's disease. *British Journal of Dermatology* 142:702-711 (2000).
4. Proby, C. M., et al. Development of chimeric molecules for recognition and targeting of antigen-specific B cells in pemphigus vulgaris. *British Journal of Dermatology* 142:321-330 (2000).
5. Wahl, J. K. III, et al. The amino- and carboxyl-terminal tails of (beta)-catenin reduce its affinity for desmoglein 2. *J Cell Sci.* 113:1737-1745 (2000).

RELATED PRODUCTS

Product	Clone	Cat. No.
Mouse anti-Desmoglein-1	27B2	32-6000
Mouse anti-Desmoglein-2	6D8	32-6100
Mouse anti-Desmoglein-3	5G11	32-6300
Mouse anti-Plakophilin-1	10B2	32-5700
Mouse anti- α -Catenin	α CAT-7A4	13-9700
Mouse anti- β -Catenin	CAT-5H10	13-8400
Mouse anti- γ -Catenin	PG-11E4	13-8500

Product	Conjugate	Cat. No.
Protein A	Sepharose [®] 4B	10-1041
rec-Protein G	Sepharose [®] 4B	10-1241

Conjugate	ZyMAX™ Goat x Mouse IgG (H+L)
Purified	81-6500
FITC	81-6511
TRITC	81-6514
Cy™ ₃	81-6515
Cy™ ₅	81-6516
HRP	81-6520
AP	81-6522
Biotin	81-6540

Zymed[®] and ZyMAX™ are trademarks of Zymed Laboratories Inc. Cy™ is a trademark of Amersham Life Sciences, Inc. Sepharose[®] is a registered trademark of Pharmacia LKB.

For Research Use Only

www.invitrogen.com

Invitrogen Corporation • 542 Flynn Rd • Camarillo • CA 93012 • Tel: 800.955.6288 • E-mail: techsupport@invitrogen.com

PI326200

(Rev 10/08) DCC-08-1089

Important Licensing Information - These products may be covered by one or more Limited Use Label Licenses (see the Invitrogen Catalog or our website, www.invitrogen.com). By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.