

Qty: 100 μ g/400 μ L

Rabbit anti-Mina53 (N-term)

Catalog No. 40-9500

Lot No.

Rabbit anti-Mina53 (N-term)

FORM

This polyclonal antibody is supplied as a 400 µL aliquot at a concentration of 0.25 mg/mL in phosphate buffered saline (pH 7.4) containing 0.1% sodium azide. This antibody is epitope-affinity purified from rabbit antiserum.

PAD: ZMD.516

IMMUNOGEN

Synthetic peptide derived from the N-terminal region of the human, mouse, rat, and bovine Mina53 proteins

SPECIFICITY

This antibody is specific for the N-terminal region of the Mina53 (Myc-induced nuclear antigen, mineral dust-induced gene (mdig)) protein. On Western blots, it identifies the target band at ~53 kDa.

REACTIVITY

Reactivity has been confirmed with human HT29, A549, Jurkat, HeLa cell and HeLa nuclear lysates. Based on amino acid sequence homology, reactivity with mouse, rat, and bovine is expected.

Sample	Western Blotting
Human	+++
Mouse	ND
Rat	ND
Bovine	ND

(Excellent +++, Good++, Poor +, No reactivity 0, Not applicable N/A, Not Determined ND)

USAGE

Working concentrations for specific applications should be determined by the investigator. Appropriate concentrations will be affected by several factors, including secondary antibody affinity, antigen concentration, sensitivity of detection method, temperature and length of incubations, etc. The suitability of this antibody for applications other than those listed below has not been determined. The following concentration ranges are recommended starting points for this product.

Western Blotting: 1-3 μg/mL

STORAGE

Store at 2-8°C for up to one month. Store at -20°C for long-term storage. Avoid repeated freezing and thawing.

(cont'd)

www.invitrogen.com

 $Invitrogen\ Corporation \bullet 542\ Flynn\ Rd \bullet Camarillo \bullet CA\ 93012 \bullet Tel:\ 800.955.6288 \bullet E-mail: \\ \underline{techsupport@invitrogen.com}$

BACKGROUND

Mina53 (Myc-induced nuclear antigen) is a gene that encodes a 53 kDa nuclear and nucleolar protein involved in cell proliferation. Specific inhibition of *Mina53* using an RNA interference method has been shown to severely suppress cell proliferation, suggesting that Mina53 contributes to cell growth and carcinogenesis induced by *c-Myc.*¹ Mina53 has been described as a potential tumor marker in colon cancer² and lung cancer,³ where expression levels are distinct between neoplastic and normal tissues. In esophageal squamous cell carcinoma (ESCC), Mina53 has been proposed as a characteristic feature, with higher expression levels linked to shorter patient survival periods.⁴

REFERENCES

- 1. Tsuneoka M, et al. J Biol Chem 277(38):35450-35459, 2002.
- 2. Teye K, et al. Am J Pathol 164(1):205-216, 2004.
- 3. Zhang Y, et al. Oncogene 24(31):4873-4882, 2005.
- 4. Tsuneoka M, et al. Clin Cancer Res 10(21):734-7356, 2004.

RELATED PRODUCTS

Product	Conjugate	Cat. No.
Protein A	Sepharose [®] 4B	10-1041
rec-Protein G	Sepharose [®] 4B	10-1241

Conjugate	ZyMAX™ Goat x Rabbit IgG (H+L)	ZyMAX™ Goat x Mouse IgG (H+L)
Purified	81-6100	81-6500
FITC	81-6111	81-6511
TRITC	81-6114	81-6514
Су™3	81-6115	81-6515
Су™5	81-6116	81-6516
HRP	81-6120	81-6520
AP	81-6122	81-6522
Biotin	81-6140	81-6540

Zymed® and ZyMAX™ are trademarks of Zymed Laboratories Inc. Cy™ and Sepharose® are trademarks of Amersham Biosciences Ltd.

For Research Use Only

LF051121