



Qty: 100 µg/400 µL

Rabbit anti-Galectin-1 (N-term)

Catalog No. 42-5900

Lot No.

Rabbit anti-Galectin-1 (N-term)

FORM

This polyclonal antibody is supplied as a 400 µL aliquot at a concentration of 0.25 mg/mL in phosphate buffered saline (pH 7.4) containing 0.1% sodium azide. This antibody is epitope-affinity purified from rabbit antiserum.

PAD: ZMD.580

IMMUNOGEN

Synthetic peptide derived from the N-terminal region of the human, mouse, rat, and pig galectin-1 proteins, which differ from bovine, sheep, and dog by two amino acid changes

SPECIFICITY

This antibody is specific for the galectin-1 (lectin, β-galactoside-binding lectin L-14-I, lactose-binding lectin 1, S-lac lectin 1, galaptin) protein. On Western blots, it identifies the target band at ~14 kDa.

REACTIVITY

Reactivity has been confirmed with human HUVEC and PC-3 and mouse NIH 3T3 cell lysates and ovary homogenates. Based on amino acid sequence homology, reactivity with bovine, sheep, pig, rat, and dog is expected.

Sample	Western Blotting
Human	+++
Mouse	+++
Bovine	ND
Sheep	ND
Pig	ND
Rat	ND
Dog	ND

(Excellent +++, Good++, Poor +, No reactivity 0, Not applicable N/A, Not Determined ND)

USAGE

Working concentrations for specific applications should be determined by the investigator. Appropriate concentrations will be affected by several factors, including secondary antibody affinity, antigen concentration, sensitivity of detection method, temperature and length of incubations, etc. The suitability of this antibody for applications other than those listed below has not been determined. The following concentration ranges are recommended starting points for this product.

Western Blotting: 1-3 µg/mL

STORAGE

Store at 2-8°C for up to one month. Store at -20°C for long-term storage. Avoid repeated freezing and thawing.

(cont'd)

www.invitrogen.com

Invitrogen Corporation • 542 Flynn Rd • Camarillo • CA 93012 • Tel: 800.955.6288 • E-mail: techsupport@invitrogen.com

PI425900

(Rev 10/08) DCC-08-1089

Important Licensing Information - These products may be covered by one or more Limited Use Label Licenses (see the Invitrogen Catalog or our website, www.invitrogen.com). By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BACKGROUND

The galectins are a family of β -galactoside-binding proteins implicated in modulating cell-cell and cell-matrix interactions. Galectin-1 is a homodimeric protein that binds Igals3bp and is widely expressed in cardiac, smooth, and skeletal muscle, neurons, thymus, kidney, placenta and hematopoietic cells. Galectin-1 has a diverse range of activities in relation to cell survival and proliferation including acting as a mitogen, inhibiting cellular proliferation, and promoting cellular apoptosis.¹ Galectin-1 binds to cell surface receptors such as CD45, CD3, CD4, CD7,³ and it induces T cell death.² Galectin-1 is one of the major proteins expressed in human mesenchymal stem cells (hMSC).³ It is secreted and found at the cell surface of MSC, participating in extracellular matrix-cell interactions. It is also expressed in a subset of slowly dividing subventricular zone astrocytes, which includes the neural stem cells.⁴ In the presence of galectin-1 human fetal mesenchymal stem cells also assume a muscular phenotype indicating that this protein may have relevance in developing therapies for muscular dystrophies.⁵ Galectin-1 can also be secreted by activated CD8(+) T-lymphocytes. This secreted protein acts as a soluble adhesion molecule by facilitating attachment of HIV-1 to the cell surface, suggesting that galectin-1 might affect the pathogenesis of HIV-1 infection.⁶ Galectin-1 is also a potential cancer target contributing to tumor evasion of immune responses.⁷

REFERENCES

1. Scott K & Weinberg C. *Glycoconj J* 19(7-9):467-477, 2004.
2. Stillman BN, et al. *J Immunol* 176(2):778-789, 2006.
3. Kadri T, et al. *Stem Cells Dev* 14(2):204-212, 2005.
4. Sakaguchi M, et al. *PNAS* 103(18):7112-7117, 2006.
5. Chan J, et al. *Stem Cells* 24(8):1879-1891, 2006.
6. Ouellet M, et al. *J Immunol* 174(7):4120-4126, 2005.
7. Rabinovich GA. *Br J Cancer* 92(7):1188-1192, 2005.

RELATED PRODUCTS

Product	Conjugate	Cat. No.
Protein A	Sepharose [®] 4B	10-1041
rec-Protein G	Sepharose [®] 4B	10-1241

Conjugate	ZyMAX[™] Goat x Rabbit IgG (H+L)	ZyMAX[™] Goat x Mouse IgG (H+L)
Purified	81-6100	81-6500
FITC	81-6111	81-6511
TRITC	81-6114	81-6514
Cy [™] 3	81-6115	81-6515
Cy [™] 5	81-6116	81-6516
HRP	81-6120	81-6520
AP	81-6122	81-6522
Biotin	81-6140	81-6540

Zymed[®] and ZyMAX[™] are trademarks of Zymed Laboratories Inc. Cy[™] and SEPHAROSE[®] are trademarks of GE Healthcare.

For Research Use Only

MZ060726

www.invitrogen.com

Invitrogen Corporation • 542 Flynn Rd • Camarillo • CA 93012 • Tel: 800.955.6288 • E-mail: techsupport@invitrogen.com

PI425900

(Rev 10/08) DCC-08-1089

Important Licensing Information - These products may be covered by one or more Limited Use Label Licenses (see the Invitrogen Catalog or our website, www.invitrogen.com). By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.