



Qty: 100 µg/400 µL

Rabbit anti-Pannexin 1 (C-term)

Catalog No. 488100

Lot No.

Rabbit anti-Pannexin 1 (C-term)

FORM

This polyclonal antibody is supplied as a 400 µL aliquot at a concentration of 0.25 mg/mL in phosphate buffered saline (pH 7.4) containing 0.1% sodium azide. This antibody is epitope-affinity purified from rabbit antiserum.

PAD: ZMD.697

IMMUNOGEN

Synthetic peptide derived from the C-terminal region of the mouse Pannexin 1 protein (Accession# NP_062355), which is identical to rat and 92% homologous to human sequence.

SPECIFICITY

This antibody is specific for the Pannexin 1 (Pax1, PX1, innexin) protein. On Western blots, it identifies the target band at ~45 kDa.

REACTIVITY

Reactivity has been confirmed with Pannexin 1 transfected C6 cell lysates by Western blotting. Based on amino acid sequence homology, reactivity with rat and human is expected. This antibody is recommended for Western Blotting. For Immunohistochemistry (IHC) application, please refer to products: Rabbit anti-Pannexin 1 (N-term) (Cat.# 487900) and Rabbit anti-Pannexin 1 (Mid) (Cat.# 488000).

Sample	Western Blotting	Immuno-cytochemistry	Immuno-histochemistry
Human	ND	ND	ND
Mouse	+++	ND	ND
Rat	ND	ND	ND

(Excellent +++, Good++, Poor +, No reactivity 0, Not applicable N/A, Not Determined ND)

USAGE

Working concentrations for specific applications should be determined by the investigator. Appropriate concentrations will be affected by several factors, including secondary antibody affinity, antigen concentration, sensitivity of detection method, temperature and length of incubations, etc. The suitability of this antibody for applications other than that listed below has not been determined. The following concentration range is the recommended starting point for this product.

Western Blotting: 1-3 µg/mL

STORAGE

Store at 2-8°C for up to one month. Store at -20°C for long-term storage. Avoid repeated freezing and thawing.

(cont'd)

www.invitrogen.com

Invitrogen Corporation • 542 Flynn Rd • Camarillo • CA 93012 • Tel: 800.955.6288 • E-mail: techsupport@invitrogen.com

PI488100

(Rev 10/08) DCC-08-1089

Important Licensing Information - These products may be covered by one or more Limited Use Label Licenses (see the Invitrogen Catalog or our website, www.invitrogen.com). By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BACKGROUND

Pannexins constitute a new family of gap junction type proteins. Besides the connexin proteins, Pannexin 1 and 2 seem to be molecular components of neuronal gap junctions ("electrical synapses") that are thought to form connections between principal cells in the hippocampus.¹ Functional expression in paired *Xenopus* oocytes indicated that pannexins are capable of forming communicating junctions.² In addition to forming gap junction channels in paired oocytes, Pannexin 1 can also form a mechanosensitive and ATP-permeable channel in the nonjunctional plasma membrane.³ The ATP-releasing Pannexin 1 hemichannels have proven to be important for cell-cell communication and signal processing.^{3,4,5} Immunohistochemistry revealed postsynaptic localization of Pannexin 1 in rodent hippocampal and cortical principal neurons accumulating at postsynaptic densities. The asymmetric synaptic distribution of Pannexin 1 suggests that it may function in neurons as non-junctional channels (pannexons) at postsynaptic sites and comprises a novel component of the postsynaptic protein complex.⁶

REFERENCES

1. Bruzzone R, et al. *Proc Natl Acad Sci USA* 100(23):13644-13649, 2003.
2. Vanden Abeele F, et al. *J Cell Biol* 174(4):535-546, 2006.
3. Huang YJ, et al. *Proc Natl Acad Sci USA* 104(15):6436-6441, 2007.
4. Romanov RA, et al. *EMBO J* 26(3):657-667, 2007.
5. Thompson RJ, et al. *Science* 312(5775):924-927, 2006.
6. Zoidl G, et al. *Neuroscience* 146(1):9-16, 2007.

RELATED PRODUCTS

Product	Conjugate	Cat. No.
Protein A	Sepharose 4B	10-1041
rec-Protein G	Sepharose 4B	10-1241
ZyMAX™ Goat anti-rabbit IgG	Unconjugated	81-6100
ZyMAX™ Goat anti-mouse IgG	Unconjugated	81-6500

Secondary antibody conjugates.

Conjugate	Goat anti-rabbit IgG (H+L)	Goat anti-mouse IgG (H+L)	Ex/Em*	Fluorescence similar to--
Alexa Fluor® 488	A11008	A11001	495/519	FITC
Alexa Fluor® 555	A21428	A21422	555/565	Cy3
Alexa Fluor® 594	A11012	A11005	590/617	Texas Red
Alexa Fluor® 647	A21244	A21235	650/668	Cy5
HRP	81-6120	81-6520	NA**	NA
AP	81-6122	81-6522	NA	NA
Biotin	B2770	B2763	NA	NA

*Excitation/emission (nm); **Not applicable

For additional secondary antibody conjugates, visit www.invitrogen.com/antibodies

For Research Use Only

www.invitrogen.com

Invitrogen Corporation • 542 Flynn Rd • Camarillo • CA 93012 • Tel: 800.955.6288 • E-mail: techsupport@invitrogen.com

PI488100

(Rev 10/08) DCC-08-1089

Important Licensing Information - These products may be covered by one or more Limited Use Label Licenses (see the Invitrogen Catalog or our website, www.invitrogen.com). By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.