



62-6540 2 ml
Biotin-Goat Anti-
Mouse IgG (H+L)
Lot No. See product label

Biotin-Goat Anti-Mouse IgG (H+L)

INVITROGEN AFFINITY PURIFIED ANTIGEN SPECIFIC ANTIBODY

FORM: Liquid in 10 mM phosphate-0.15 M NaCl buffer, pH 7.4, containing 1% bovine serum albumin, 0.1% sodium azide, and 50% glycerol. The antibody concentration is 0.75 mg/ml.

AFFINITY PURIFIED GOAT ANTI-MOUSE IgG (H+L) SPECIFIC ANTIBODY is biotinylated under optimum conditions and is free of unconjugated biotin. The antibody has been absorbed with solid phase human serum protein.

SUGGESTED WORKING DILUTION FOR EIA: 1:8,000 to 1:10,000 with phosphate buffered saline containing 0.05% Tween 20. However, working dilution for specific application should be determined by the investigator to determined the best conditions.

STORAGE: Store at 4°C.

REMARKS: For research use only. Not for human or drug use.

Research Use Only

www.invitrogen.com

Invitrogen Corporation • 542 Flynn Rd • Camarillo • CA 93012 • Tel: 800.955.6288

E-mail: techsupport@invitrogen.com

PI626540

(Rev 10/08) DCC-08-1089

Important Licensing Information - These products may be covered by one or more Limited Use Label Licenses (see the Invitrogen Catalog or our website, www.invitrogen.com). By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.



62-6540 2 ml
Biotin-Goat Anti-
Mouse IgG (H+L)
Lot No. See product label

Biotin-Goat Anti-Mouse IgG (H+L)

INVITROGEN AFFINITY PURIFIED ANTIGEN SPECIFIC ANTIBODY

FORM: Liquid in 10 mM phosphate-0.15 M NaCl buffer, pH 7.4, containing 1% bovine serum albumin, 0.1% sodium azide, and 50% glycerol. The antibody concentration is 0.75 mg/ml.

AFFINITY PURIFIED GOAT ANTI-MOUSE IgG (H+L) SPECIFIC ANTIBODY is biotinylated under optimum conditions and is free of unconjugated biotin. The antibody has been absorbed with solid phase human serum protein.

SUGGESTED WORKING DILUTION FOR EIA: 1:8,000 to 1:10,000 with phosphate buffered saline containing 0.05% Tween 20. However, working dilution for specific application should be determined by the investigator to determined the best conditions.

STORAGE: Store at 4°C.

REMARKS: For research use only. Not for human or drug use.

Research Use Only

www.invitrogen.com

Invitrogen Corporation • 542 Flynn Rd • Camarillo • CA 93012 • Tel: 800.955.6288

E-mail: techsupport@invitrogen.com

PI626540

(Rev 10/08) DCC-08-1089

Important Licensing Information - These products may be covered by one or more Limited Use Label Licenses (see the Invitrogen Catalog or our website, www.invitrogen.com). By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.