



Qty: 100 µg/400 µL

Rabbit anti-Claudin-1

Catalog No. 51-9000

Lot No. See product label

Rabbit anti-Claudin-1

FORM

This polyclonal antibody is supplied as a 400 µL aliquot at a concentration of 0.25 mg/mL in phosphate buffered saline (pH 7.4) containing 0.1% sodium azide. The antibody is epitope-affinity-purified from rabbit antiserum.

POLYCLONAL ANTIBODY DESIGNATION (PAD): JAY.8

ISOTYPE: Rabbit Ig

IMMUNOGEN

Synthetic peptide derived from the C-terminus of the human Claudin-1 protein

SPECIFICITY

This antibody is specific for the ~ 22 kDa Claudin-1 protein, and does not cross-react with Claudin-3.

REACTIVITY

Reactivity has been confirmed with human Caco-2 and canine MDCK cell lysates and rat kidney homogenates by Western blotting and immunofluorescence, and with human skin tissue by immunohistochemistry.

Sample	ELISA	Immuno-fluorescence	Western blotting	Immuno-histochemistry (FFPE)*
Dog		not tested	+	not tested
Human		not tested	+	+
Rat		+	not tested	not tested
Immunogen	+	not tested	not tested	not tested

USAGE

Working concentrations for specific applications should be determined by the investigator. Appropriate concentrations will be affected by several factors, including secondary antibody affinity, antigen concentration, sensitivity of detection method, temperature and length of incubations, etc. The suitability of this antibody for applications other than those listed below has not been determined. The following concentration ranges are recommended starting points for this product.

ELISA: 0.1-1.0 µg/mL
Western Blotting: 1-2 µg/mL
Immunofluorescence: 5-10 µg/mL
Immunohistochemistry*: 5-10 µg/mL

* For best results in immunohistochemistry with formalin-fixed, paraffin-embedded tissues, heat induced epitope retrieval (HIER) with EDTA, pH 8.0, is required prior to staining.

STORAGE

Store at 2-8°C for up to one month. Store at -20°C for long-term storage. Avoid repeated freezing and thawing.

(cont'd)

www.invitrogen.com

Invitrogen Corporation • 542 Flynn Rd • Camarillo • CA 93012 • Tel: 800.955.6288 • E-mail: techsupport@invitrogen.com

PI519000

(Rev 10/08) DCC-08-1089

Important Licensing Information - These products may be covered by one or more Limited Use Label Licenses (see the Invitrogen Catalog or our website, www.invitrogen.com). By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BACKGROUND

Tight junctions are specialized regions of cell-cell contact that are particularly abundant in luminal epithelial cell sheets. In freeze-fracture electron micrographs, tight junctions are visualized as belt-like bands of anastomosing sealing strands (TJ strands) that completely encircle the lateral surfaces of each cell. TJ strands on adjacent cells interact with each other to form a "molecular gasket" that prevents ions, water and other molecules from leaking between cells and thus, from one side of the sheet to the other. In addition to this "barrier" function, the "fence" function of tight junctions plays an important role in maintaining epithelial cell-polarity by blocking the diffusion of membrane proteins between apical (luminal) and basolateral cell surfaces.

Several peripheral membrane proteins are associated with tight junctions including ZO-1, ZO-2, ZO-3, cingulin, the 7H6 antigen, Rab-3b, and symplekin.¹⁻⁶ Suggested roles for these proteins include involvement in tight junction assembly and maintenance, signal transduction, and the regulation of tight junction permeability. A growing body of evidence suggests that actin filaments play a major role in regulating tight junction permeability.

Until recently, the only transmembrane protein known to be associated with tight junctions was occludin, an ~65 kDa protein with four transmembrane domains. Despite widespread expectation, a critical structural role for occludin in TJ strands was ruled out by the observation of apparently normal tight junctions formed between cells disrupted at both occludin alleles.⁷ A closer examination of isolated tight junctions uncovered two related ~22 kDa, four-transmembrane domain proteins, claudin-1 and claudin-2, with no similarity to occludin. In contrast to occludin, which induces only a small number of short strands at cell-cell contact sites when introduced into fibroblasts lacking tight junctions, claudin-1 and -2 induce networks of strands characteristic of true tight junctions.^{8,9} Though inconclusive, these findings suggest that claudin-1 and -2 are major structural components of TJ strands and that occludin plays some other accessory role. Excitement in the tight junction field continues to rise following the recent discovery of claudins -3, -4, -5, -6, -7, and -8 and experiments suggesting that tight junctions in different tissues are comprised of different sets of claudin family proteins.¹⁰

REFERENCES

1. Anderson JM. *Curr Biol* 6:382-384, 1996.
2. Anderson JM, Van Itallie CM. *Am J Physiol* 269:G467-475, 1995.
3. Howarth AG, Stevenson BR. *Adv Struct Biol* 4:25-39, 1996.
4. Stevenson BR, Keon BH. *Ann Rev Cell Dev Biol* 14:89-109, 1998.
5. Tsukita S, et al. *Cell Struct Funct* 21:381-385, 1996.
6. Yap AS, et al. *Mem Biol* 163:159-167, 1998.
7. Saitou M, et al. *J Cell Biol* 141:397-408, 1998.
8. Furuse M, et al. *J Cell Biol* 143:391-401, 1998.
9. Tsukita S, Furuse M. *Genes Cells* 3:569-573, 1998.
10. Morita K, et al. *PNAS* 96:511-516, 1999.

RELATED PRODUCTS

Product	Clone/PAD*	Cat. No.
Rb x Claudin-2	MH44	51-6100
Ms x ZO-1	ZO1-1A12	33-9100
Rb x ZO-1	Z-R1	61-7300
Rb x ZO-2	--	71-1400
Ms x Occludin	OC-3F10	33-1500
Rb x Occludin	Z-T22	71-1500

PAD: Polyclonal Antibody Designation

Conjugate	ZyMAX™ Goat x Rabbit IgG (H+L)	ZyMAX™ Goat x Mouse IgG (H+L)
Purified	81-6100	81-6500
FITC	81-6111	81-6511
TRITC	81-6114	81-6514
Cy™3	81-6115	81-6515
Cy™5	81-6116	81-6516
HRP	81-6120	81-6520
AP	81-6122	81-6522
Biotin	81-6140	81-6540

Zymed® and ZyMAX™ are trademarks of Zymed Laboratories Inc. Cy™ is a trademark of Amersham Life Sciences, Inc.

For Research Use Only

Leej010216

www.invitrogen.com

Invitrogen Corporation • 542 Flynn Rd • Camarillo • CA 93012 • Tel: 800.955.6288 • E-mail: techsupport@invitrogen.com

PI519000

(Rev 10/08) DCC-08-1089

Important Licensing Information - These products may be covered by one or more Limited Use Label Licenses (see the Invitrogen Catalog or our website, www.invitrogen.com). By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.