



65-6140 2 ml
Biotin-Goat Anti-
Rabbit IgG (H+L)

BIOTINYLATED-GOAT ANTI-RABBIT IgG (H+L)

INVITROGEN AFFINITY PURIFIED ANTIGEN SPECIFIC ANTIBODY *DOUBLE STAINING GRADE*

FORM: Liquid in 10 mM phosphate-0.15 M NaCl buffer, pH 7.4, containing 1% bovine serum albumin, 0.1% sodium azide and 50% glycerol. The antibody concentration is 0.60 mg/ml.

AFFINITY PURIFIED GOAT ANTI-RABBIT IgG (H+L) SPECIFIC ANTIBODY* is biotinylated under optimum conditions and is free of unconjugated biotin. This antibody had been absorbed and tested to show minimal cross-reactivity to the serum proteins of the following species: mouse, human, and rat.

SUGGESTED WORKING DILUTION FOR EIA: 1:8,000 to 1:10,000 with phosphate buffered saline containing 0.05% Tween 20. However, working dilution for specific application should be determined by the investigator to obtain the best conditions.

STORAGE: Store at 2-8°C . Avoid repeated freezing and thawing.

REMARKS: Product contains 0.1% sodium azide as preservative. This product has not been assayed for species cross-reactivity unless otherwise indicated.

For research use only.

CAUTION: Not intended for human or animal therapeutic or diagnostic use.

www.invitrogen.com

Invitrogen Corporation • 542 Flynn Rd • Camarillo • CA 93012 • Tel: 800.955.6288

E-mail: techsupport@invitrogen.com

PI656140

(Rev 12/09) DCC-09-1970

Important Licensing Information - These products may be covered by one or more Limited Use Label Licenses (see the Invitrogen Catalog or our website, www.invitrogen.com). By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.



65-6140 2 ml
Biotin-Goat Anti-
Rabbit IgG (H+L)

BIOTINYLATED-GOAT ANTI-RABBIT IgG (H+L)

INVITROGEN AFFINITY PURIFIED ANTIGEN SPECIFIC ANTIBODY *DOUBLE STAINING GRADE*

FORM: Liquid in 10 mM phosphate-0.15 M NaCl buffer, pH 7.4, containing 1% bovine serum albumin, 0.1% sodium azide and 50% glycerol. The antibody concentration is 0.60 mg/ml.

AFFINITY PURIFIED GOAT ANTI-RABBIT IgG (H+L) SPECIFIC ANTIBODY* is biotinylated under optimum conditions and is free of unconjugated biotin. This antibody had been absorbed and tested to show minimal cross-reactivity to the serum proteins of the following species: mouse, human, and rat.

SUGGESTED WORKING DILUTION FOR EIA: 1:8,000 to 1:10,000 with phosphate buffered saline containing 0.05% Tween 20. However, working dilution for specific application should be determined by the investigator to obtain the best conditions.

STORAGE: Store at 2-8°C. Avoid repeated freezing and thawing.

REMARKS: Product contains 0.1% sodium azide as preservative. This product has not been assayed for species cross-reactivity unless otherwise indicated.

For research use only.

CAUTION: Not intended for human or animal therapeutic or diagnostic use.

www.invitrogen.com

Invitrogen Corporation • 542 Flynn Rd • Camarillo • CA 93012 • Tel: 800.955.6288

E-mail: techsupport@invitrogen.com

PI656140

(Rev 12/09) DCC-09-1970

Important Licensing Information - These products may be covered by one or more Limited Use Label Licenses (see the Invitrogen Catalog or our website, www.invitrogen.com). By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.